

Minimal time for the continuity equation controlled by a localized perturbation of the velocity vector field

Michel Duprez*

Morgan Morancey[†]

Francesco Rossi[‡]

November 28, 2019

Abstract

In this work, we study the minimal time to steer a given crowd to a desired configuration. The control is a vector field, representing a perturbation of the crowd velocity, localized on a fixed control set. We will assume that there is no interaction between the agents.

We give a characterization of the minimal time both for microscopic and macroscopic descriptions of a crowd. We show that the minimal time to steer one initial configuration to another is related to the condition of having enough mass in the control region to feed the desired final configuration.

The construction of the control is explicit, providing a numerical algorithm for computing it. We finally give some numerical simulations.

1 Introduction and main results

In recent years, the study of systems describing a crowd of interacting autonomous agents has drawn a great interest from the control community. A better understanding of such interaction phenomena can have a strong impact in several key applications, such as road traffic and egress problems for pedestrians. For a few reviews about this topic, see *e.g.* [5, 6, 12, 22, 29, 31, 39, 47].

Beside the description of interactions, it is now relevant to study problems of control of crowds, *i.e.* of controlling such systems by acting on few agents, or on a small subset of the configuration space. Roughly speaking, basic problems for such models are controllability (*i.e.* reachability of a desired configuration) and optimal control (*i.e.* the minimization of a given functional). We already addressed the controllability problem in [24], thus identifying reachable configurations for crowd models. The present article deals with the subsequent step, that is the study of a classical optimal control problem: the minimal time to reach a desired configuration.

The nature of the control problem relies on the model used to describe the crowd. Two main classes are widely used. In microscopic models, the position of each agent is clearly identified; the crowd dynamics is described by an ordinary differential equation of large dimension, in which coupling terms represent interactions between agents. For control of such models, a large literature is available, see *e.g.* reviews [10, 35, 36], as well as applications, both to pedestrian crowds [26, 38] and to road traffic [13, 28].

In macroscopic models, instead, the idea is to represent the crowd by the spatial density of agents; in this setting, the evolution in time of the density solves a partial differential equation, usually of transport type. Nonlocal terms (such as convolutions) model interactions between agents. To our knowledge, there exist few studies of control of this family of equations. In [43], the authors provide approximate alignment

*CEREMADE, Université Paris-Dauphine & CNRS UMR 7534, Université PSL, 75016 Paris, France. (mduprez@math.cnrs.fr).

[†]Aix Marseille Université, CNRS, Centrale Marseille, I2M, Marseille, France. (morgan.morancey@univ-amu.fr).

[‡]Dipartimento di Matematica “Tullio Levi-Civita”, Università degli Studi di Padova, Via Trieste 63, 35121 Padova, Italy. (francesco.rossi@math.unipd.it).

of a crowd described by the macroscopic Cucker-Smale model [23]. The control is the acceleration, and it is localized in a control region ω which moves in time. In a similar situation, a stabilization strategy has been established in [14, 15], by generalizing the Jurdjevic-Quinn method to partial differential equations. A different approach is given by mean-field type control, *i.e.* control of mean-field equations, and by mean-field games modelling crowds, see *e.g.* [1, 3, 7, 8, 16, 27]. In this case, problems are often of optimization nature, *i.e.* the goal is to find a control minimizing a given cost, with no final constraint. A notable exception is the so-called planning problem, both in the pure transport setting [40, 32] and with an additional diffusion term [2, 45, 46]. In this article, there is no individual optimization, thus these results do not apply to our setting.

This article deals with the problem of steering one initial configuration to a final one in minimal time. As stated above, we recently discussed in [24] the problem of controllability for the systems described here, which main results are recalled in Section 2.3. We proved that one can approximately steer an initial to a final configuration if they satisfy the Geometric Condition 1 recalled below. Roughly speaking, it requires that the whole initial configuration crosses the control set ω forward in time, and the final one crosses it backward in time. From now on, we will always assume that this condition is satisfied, so to ensure that the final configuration is approximately reachable.

When the controllability property is satisfied, it is then interesting to study minimal time problems. Indeed, from the theoretical point of view, it is the first problem in which optimality conditions can be naturally defined. More related to applications described above, minimal time problems play a crucial role: egress problems can be described in this setting, while traffic control is often described in terms of minimization of (maximal or average) total travel time.

Notice that the minimal time issue in a control problem can be a consequence of various unrelated phenomena. For instance, imposing constraints on the control or on the state can lead to a positive minimal time even if the considered evolution equation enjoys an infinite propagation speed (see for instance [37]). On the contrary, considering evolution equations with a finite speed of propagation, such as transport or wave equations, naturally imply a minimal null control time when the control is localized. Our contribution fits in the second setting.

For microscopic models, the dynamics can be written in terms of finite-dimensional control systems. For this reason, minimal time problems can sometimes be addressed with classical (linear or non-linear) control theory, see *e.g.* [4, 9, 33, 48]. Instead, very few results are known for macroscopic models, that can be recasted in terms of control of the transport equation. The linear case is classical, see *e.g.* [21]. Instead, more recent developments in the non-linear case (based on generalization of differential inclusions) have been recently described in [17, 18, 19].

The originality of our research lies in the constraint given on the control: it is a perturbation of the velocity vector field localized in a given region ω of the space. Such constraint is highly non-trivial, since the control problem is clearly non-linear even though the uncontrolled dynamics is. To the best of our knowledge, minimal time problems with this constraints have not been studied, neither for microscopic nor for macroscopic models. We first study a microscopic model, where the crowd is described by a vector with nd components ($n, d \in \mathbb{N}^*$) representing the positions of n agents in \mathbb{R}^d . The natural (uncontrolled) vector field is denoted by $v : \mathbb{R}^d \rightarrow \mathbb{R}^d$, assumed Lipschitz and uniformly bounded. We act on the vector field in a fixed subdomain ω of the space, which will be a bounded open connected subset of \mathbb{R}^d . The admissible controls are thus functions of the form $\mathbf{1}_\omega u : \mathbb{R}^d \times \mathbb{R}^+ \rightarrow \mathbb{R}^d$. The control models an external action (*e.g.* dynamic signs for road traffic, or traffic agents in pedestrian crowds) located in a fixed area, that interacts with all agents in such given area. Since ω is fixed, it can act on predefined locations only.

We then consider the following ordinary differential equation (microscopic model)

$$\begin{cases} \dot{x}_i(t) = v(x_i(t)) + \mathbf{1}_\omega(x_i(t))u(x_i(t), t), \\ x_i(0) = x_i^0 \end{cases} \quad i = 1, \dots, n, \quad (1)$$

where $X^0 := \{x_1^0, \dots, x_n^0\}$ is the initial configuration of the crowd.

We also study a macroscopic model, where the crowd is represented by its density, that is a time-evolving probability measure $\mu(t)$ defined on the space \mathbb{R}^d . We consider the same natural vector field v , control region ω and admissible controls $\mathbb{1}_\omega u$. The dynamics is given by the following linear transport equation (macroscopic model)

$$\begin{cases} \partial_t \mu + \nabla \cdot ((v + \mathbb{1}_\omega u)\mu) = 0 & \text{in } \mathbb{R}^d \times \mathbb{R}^+, \\ \mu(\cdot, 0) = \mu^0 & \text{in } \mathbb{R}^d, \end{cases} \quad (2)$$

where μ^0 is the initial density of the crowd. The function $v + \mathbb{1}_\omega u$ represents the vector field acting on μ . To a microscopic configuration $X := \{x_1, \dots, x_n\}$, we can associate the empirical measure

$$\mu := \sum_{i=1}^n \frac{1}{n} \delta_{x_i}. \quad (3)$$

With this notation, System (1) is a particular case of System (2). This identification can be applied only if the different microscopic agents are considered identical or interchangeable, as it is often the case for crowd models with a large number of agents. This identification will be used several times in the following, namely to approximate macroscopic models with microscopic ones.

Systems (1) and (2) are first approximations for crowd modeling, since the uncontrolled vector field v is given, and it does not describe interactions between agents. Nevertheless, it is necessary to understand control properties for such simple equations as a first step, before dealing with vector fields depending on the crowd itself. Thus, in a future work, using the results for linear equations presented here, we will study control problems for crowd models with a non-local term $v[x_1, \dots, x_n]$ in the microscopic model (1) and $v[\mu]$ in the the macroscopic model (2).

From now on, we always assume that the following geometric condition is satisfied:

Condition 1 (Geometric condition). Let μ^0, μ^1 be two probability measures on \mathbb{R}^d satisfying:

- (i) For each $x^0 \in \text{supp}(\mu^0)$, there exists $t^0 > 0$ such that $\Phi_{t^0}^v(x^0) \in \omega$, where Φ_t^v is the *flow* associated to v (see Definition 2.3 below).
- (ii) For each $x^1 \in \text{supp}(\mu^1)$, there exists $t^1 > 0$ such that $\Phi_{-t^1}^v(x^1) \in \omega$.

The Geometric Condition 1 means that each particle crosses the control region. It is illustrated in Figure 1. It is the minimal condition that we can expect to steer any initial condition to any target. Indeed, if Item (i) of the Geometric Condition 1 is not satisfied, then there exists a whole subpopulation of the measure μ^0 that never intersects the control region, thus the localized control cannot act on it.




Figure 1: Geometric condition.

To ensure the well-posedness of Systems (1) and (2), we fix regularity conditions on the natural vector field v and the control $\mathbb{1}_\omega u$ as follows:

Condition 2 (Carathéodory condition). Let v and $\mathbb{1}_\omega u$ be such that

- (i) The applications $x \mapsto v(x)$ and $x \mapsto \mathbb{1}_\omega u(x, t)$ for each $t \in \mathbb{R}$ are Lipschitz.
- (ii) For all $x \in \mathbb{R}^d$, the application $t \mapsto \mathbb{1}_\omega u(x, t)$ is measurable.
- (iii) There exists $M > 0$ such that $\|v\|_\infty \leq M$ and $\|\mathbb{1}_\omega u\|_\infty \leq M$.

For arbitrary macroscopic models, one can expect approximate controllability only, since for general measures there exists no homeomorphism sending one to another. Indeed, if we impose the Carathéodory condition, then the flow $\Phi_t^{v+\mathbb{1}_\omega u}$ is an homeomorphism (see [9, Th. 2.1.1]). For more details, see Remark 2.

Similarly, in the microscopic case, the control vector field $\mathbb{1}_\omega u$ cannot separate points, due to uniqueness of the solution of System (1). In the microscopic model, we then assume that the initial configuration X^0 and the final one X^1 are disjoint, in the following sense:

Definition 1.1. A configuration $X = \{x_1, \dots, x_n\}$ is disjoint if $x_i \neq x_j$ for all $i \neq j$.

In other words, in a disjoint configuration, two agents cannot lie in the same point of the space. Since we deal with velocities $v + \mathbb{1}_\omega u$ satisfying the Carathéodory condition, if X^0 is a disjoint configuration, then the solution $X(t)$ to microscopic System (1) is also a disjoint configuration at each time $t \geq 0$.

In this article, we aim to study the minimal time problem. We denote by T_a the minimal time to approximately steer the initial configuration μ^0 to a final one μ^1 in the following sense: it is the infimum of times for which there exists a control $\mathbb{1}_\omega u : \mathbb{R}^d \times \mathbb{R}^+ \rightarrow \mathbb{R}^d$ satisfying the Carathéodory condition steering μ^0 arbitrarily close to μ^1 . We similarly define the minimal time T_e to exactly steer the initial configuration μ^0 to a final one μ^1 . A precise definition will be given below. Since the minimal time is not always reached, we will speak about *infimum time*.

In the sequel, we will use the following notation for all $x \in \mathbb{R}^d$:

$$\begin{aligned} \bar{t}^0(x) &:= \inf\{t \in \mathbb{R}^+ : \Phi_t^v(x) \in \bar{\omega}\}, & \bar{t}^1(x) &:= \inf\{t \in \mathbb{R}^+ : \Phi_{-t}^v(x) \in \bar{\omega}\}, \\ t^0(x) &:= \inf\{t \in \mathbb{R}^+ : \Phi_t^v(x) \in \omega\}, & t^1(x) &:= \inf\{t \in \mathbb{R}^+ : \Phi_{-t}^v(x) \in \omega\}. \end{aligned} \quad (4)$$

The quantity $\bar{t}^0(x)$ is the infimum time at which the particle localized at point x with the velocity v belongs to $\bar{\omega}$. Idem for the other quantities. Under the Geometric Condition 1, $\bar{t}^0(x)$ and $t^0(x)$ are finite for all $x \in \text{supp}(\mu^0)$, and similarly for $\bar{t}^1(x)$ and $t^1(x)$ when $x \in \text{supp}(\mu^1)$. Moreover, it is clear that it always holds $\bar{t}^j(x) \leq t^j(x)$. In some situations, this inequality can be strict. For example, in Figure 2, it holds $\bar{t}^1(x_1^1) < t^1(x_1^1)$. It is also important to remark that these quantities are not continuous with respect to x . Indeed, for instance, in Figure 2, if we shift x_1^1 to the right (resp. to the left), we observe a discontinuity of \bar{t}^1 (resp. t^1) at the point x_1^1 .

If the set ω admits everywhere an outer normal vector n , *e.g.* when it is \mathcal{C}^1 , one can have $\bar{t}^1(x_1^1) < t^1(x_1^1)$ only if $v(\Phi_{-\bar{t}^1(x_1^1)}^v(x_1^1)) \cdot n = 0$. This means that the trajectory $\Phi_{-t}^v(x_1^1)$ touches the boundary of ω with a parallel tangent vector. The proof of such simple statement can be recovered by applying the implicit function theorem.

1.1 Infimum time for microscopic models

In this section, we state the two main results about the infimum time for microscopic models. We evaluate the minimal time both for exact and approximate controllability, highlighting the different role of $t^1(x)$ and $\bar{t}^1(x)$ in the two cases.

For simplicity, we denote by

$$\bar{t}_i^0 := \bar{t}^0(x_i^0), \quad \bar{t}_i^1 := \bar{t}^1(x_i^1), \quad t_i^0 := t^0(x_i^0), \quad t_i^1 := t^1(x_i^1), \quad (5)$$




Figure 2: Example of difference between $t^1(x_1^1)$ and $\tilde{t}^1(x_1^1)$.

for $i \in \{1, \dots, n\}$. We then define

$$M_e^*(X^0, X^1) := \max\{t_i^0, t_i^1 : i = 1, \dots, n\}, \quad M_a^*(X^0, X^1) := \max\{t_i^0, \tilde{t}_i^1 : i = 1, \dots, n\}.$$

The value $M_e^*(X^0, X^1)$ has the following meaning for exact controllability: a time $T > M_e^*(X^0, X^1)$ is sufficiently large for each particle x_i^0 to enter ω and for each particle x_i^1 to enter it backward in time. The value $M_a^*(X^0, X^1)$ plays the same role for approximate controllability.

We now state our main result on the infimum time about exact control of System (1).

Theorem 1.2 (Infimum time for exact control of microscopic models). *Let $X^0 := \{x_1^0, \dots, x_n^0\}$ and $X^1 := \{x_1^1, \dots, x_n^1\}$ be two disjoint configurations (see Definition 1.1). Assume that ω is a bounded open connected set, the empirical measures (3) associated to X^0 and X^1 satisfy the Geometric Condition 1 and the velocities v and $\mathbb{1}_\omega u$ satisfy the Carathéodory Condition 2.*

Arrange the sequences $\{t_i^0\}_i$ and $\{t_j^1\}_j$ to be non-decreasingly and non-increasingly ordered, respectively. Then

$$M_e(X^0, X^1) := \max_{i \in \{1, \dots, n\}} |t_i^0 + t_i^1| \tag{6}$$

is the infimum time $T_e(X^0, X^1)$ for exact control of System (1) in the following sense:

- (i) *For each $T > M_e(X^0, X^1)$, System (1) is exactly controllable from X^0 to X^1 at time T , i.e. there exists a control $\mathbb{1}_\omega u : \mathbb{R}^d \times \mathbb{R}^+ \rightarrow \mathbb{R}^d$ steering X^0 exactly to X^1 .*
- (ii) *For each $T \in (M_e^*(X^0, X^1), M_e(X^0, X^1)]$, System (1) is not exactly controllable from X^0 to X^1 .*
- (iii) *There exists at most a finite number of times $T \in [0, M_e^*(X^0, X^1)]$ for which System (1) is exactly controllable from X^0 to X^1 .*

We give a proof of Theorem 1.2 in Section 3. As stated in item (iii), it can exist some times $T \leq M_e^*(X^0, X^1)$ at which it is possible to steer X^0 to X^1 , but it will be not entirely thanks to the control. We give an example of this situation in Remark 5 below. We now give a characterization of the minimal time for approximate control of microscopic models.

Theorem 1.3 (Infimum time for approximate control of microscopic models). *Let $X^0 := \{x_1^0, \dots, x_n^0\}$ and $X^1 := \{x_1^1, \dots, x_n^1\}$ be two disjoint configurations (see Definition 1.1). Assume that ω is a bounded open connected set, the empirical measures (3) associated to X^0 and X^1 satisfy the Geometric Condition 1 and the velocities v and $\mathbb{1}_\omega u$ satisfy the Carathéodory Condition 2. Arrange the sequences $\{t_i^0\}_i$ and $\{\tilde{t}_j^1\}_j$ to be non-decreasingly and non-increasingly ordered, respectively. Then*

$$M_a(X^0, X^1) := \max_{i \in \{1, \dots, n\}} |t_i^0 + \tilde{t}_i^1|$$

is the infimum time $T_a(X^0, X^1)$ for approximate control of System (1) in the following sense:

- (i) For each $T > M_a(X^0, X^1)$, System (1) is approximately controllable from X^0 to X^1 at time T , i.e. there exists a permutation $\sigma \in \mathbb{S}_n$ and a sequence of controls $\{\mathbf{1}_\omega u_k\}_{k \in \mathbb{N}^*}$ such that the associated solution $x_{i,k}(t)$ to System (1) satisfies $x_{i,k}(T) \xrightarrow[k \rightarrow \infty]{} x_{\sigma(i)}^1$.
- (ii) For each $T \in (M_a^*(X^0, X^1), M_a(X^0, X^1)]$, System (1) is not approximately controllable from X^0 to X^1 .
- (iii) There exists at most a finite number of times $T \in [0, M_a^*(X^0, X^1)]$ for which System (1) is approximately controllable from X^0 to X^1 .

We give a proof of Theorem 1.3 in Section 3. As for the exact controllability, one might have approximate controllability for a time $T \leq M_a^*(X^0, X^1)$, but not entirely due to the control. We refer again to Remark 5 below for an example.

It is well known that the notions of approximate and exact controllability are equivalent for finite dimensional linear systems when the control acts linearly, see *e.g.* [21]. This is not the case for System (1), which highlights the fact that we are dealing with a non-linear control problem. The difference is indeed related to the fact that for exact and approximate controllability, tangent trajectories give different behaviors. For example, in Figure 2, if we denote by $X^0 := \{x_1^0\}$ and $X^1 := \{x_1^1\}$, then it holds $M_a(X^0, X^1) < M_e(X^0, X^1)$ due to the presence of a tangent trajectory. A trajectory achieving approximate controllability is represented as dashed lines in the case $T \in (M_a(X^0, X^1), M_e(X^0, X^1))$ in Figure 2.

1.2 Infimum time for macroscopic models

In this section, we state the main result about the infimum time for approximate control of macroscopic models.

Let μ^0 and μ^1 be two probability measures, with compact support. We introduce the maps \mathcal{F}_{μ^0} and \mathcal{B}_{μ^1} defined for all $t \geq 0$ by

$$\begin{cases} \mathcal{F}_{\mu^0}(t) := \mu^0(\{x \in \text{supp}(\mu^0) : t^0(x) \leq t\}), \\ \mathcal{B}_{\mu^1}(t) := \mu^1(\{x \in \text{supp}(\mu^1) : t^1(x) \leq t\}). \end{cases}$$

The function \mathcal{F}_{μ^0} (resp. \mathcal{B}_{μ^1}) gives the quantity of mass coming from μ^0 forward in time (resp. the quantity of mass coming from μ^1 backward in time) which has entered in ω at time t . Observe that we do not decrease \mathcal{F}_{μ^0} when the mass eventually leaves ω , and similarly for \mathcal{B}_{μ^1} . Define the generalised inverse functions $\mathcal{F}_{\mu^0}^{-1}$ and $\mathcal{B}_{\mu^1}^{-1}$ of \mathcal{F}_{μ^0} and \mathcal{B}_{μ^1} given for all $m \in [0, 1]$ by

$$\begin{cases} \mathcal{F}_{\mu^0}^{-1}(m) := \inf\{t \geq 0 : \mathcal{F}_{\mu^0}(t) \geq m\}, \\ \mathcal{B}_{\mu^1}^{-1}(m) := \inf\{t \geq 0 : \mathcal{B}_{\mu^1}(t) \geq m\}. \end{cases}$$

The function $\mathcal{F}_{\mu^0}^{-1}$ is increasing, lower semi-continuous and gives the time at which a mass m has entered in ω ; similarly for $\mathcal{B}_{\mu^1}^{-1}$. Define

$$S^*(\mu^0, \mu^1) := \sup\{t^l(x) : x \in \text{supp}(\mu^l) \text{ and } l \in \{0, 1\}\}. \quad (7)$$

We then have the following main result about infimum time in the macroscopic case:

Theorem 1.4 (Infimum time for approximate control of macroscopic models). *Let μ^0 and μ^1 be two probability measures, with compact support, absolutely continuous with respect to the Lebesgue measure and satisfying the Geometric Condition 1. Assume that ω is a bounded open connected set and the velocities v and $\mathbf{1}_\omega u$ satisfy the Carathéodory Condition 2. Then*

$$S(\mu^0, \mu^1) := \sup_{m \in [0, 1]} \{\mathcal{F}_{\mu^0}^{-1}(m) + \mathcal{B}_{\mu^1}^{-1}(1 - m)\} \quad (8)$$

is the infimum time $T_a(\mu^0, \mu^1)$ to approximately steer μ^0 to μ^1 in the following sense:

(i) Assume that there exists $\alpha > 0$ for which it holds

$$\{x \in \mathbb{R}^d : d(x, \omega^c) > \varepsilon\} \text{ is connected for each } \varepsilon \in (0, \alpha), \quad (9)$$

(e.g. when ω is convex.) Then for all $T > S(\mu^0, \mu^1)$, System (2) is approximately controllable from μ^0 to μ^1 at time T .

(ii) Assume that the boundary of the evolved set

$$\omega^t := \cup_{\tau \in (0, t)} \Phi_\tau^v(\omega) \quad (10)$$

with respect to the flow Φ_τ^v of the vector field v has zero Lebesgue measure for each $t > 0$. Then, for all $T \in (S^*(\mu^0, \mu^1), S(\mu^0, \mu^1)]$, System (2) is not approximately controllable from μ^0 to μ^1 .

We give a proof of Theorem 1.4 in Section 4. We observe that the quantity $S(\mu^0, \mu^1)$ given in (8) is the continuous equivalent of $M_e(X^0, X^1)$ in (6). Contrarily to the microscopic case, System (2) can be approximately controllable at each time $T \in (0, S^*(\mu^0, \mu^1))$. We give some examples in Remark 9 below. We do not analyze the infimum time to exactly control System (2) since it is not exactly controllable with controls satisfying the Carathéodory condition. For more details, we refer to Remark 2 below.

Remark 1. The particular condition on ω^t in statement (ii) in Theorem 1.4 is crucial to ensure that AC-measures do not concentrate on the boundary. In connection with this condition, we now state the following useful Lemma, which proof is postponed to the Appendix.

It gives simple conditions, though not optimal, ensuring that the assumptions of Theorem 1.4 (ii) hold.

Lemma 1.5. *Let v be a C^1 vector field and ω an open bounded set satisfying the uniform interior cone condition (e.g. convex) as recalled in Definition A.1. Then, for each $t > 0$ the evolved set ω^t defined in (10) has boundary with zero Lebesgue measure.*

In a future work, we aim to investigate if such statement holds with less regularity of the vector field v and of the set ω . For example, we will show that the fact that ω has a boundary of zero measure does not imply that ω^t satisfies the same property.

This paper is organised as follows. In Section 2, we recall basic properties of the Wasserstein distance, ordinary differential equations and continuity equations. We prove our main results Theorems 1.2 and 1.3 in Section 3 and Theorem 1.4 in Section 4. We finally introduce an algorithm to compute the infimum time for approximate control of macroscopic models and give some numerical examples in Section 5.

2 Wasserstein distance, models and controllability

In this section, we highlight the connections of the microscopic model (1) and the macroscopic model (2) with the Wasserstein distance, that is the natural distance associated to these dynamics. We also recall our previous results obtained in [24] about controllability of these systems.

2.1 Wasserstein distance

From now on, we denote by $\mathcal{P}_c(\mathbb{R}^d)$ the space of probability measures on \mathbb{R}^d with compact support. We also denote by “AC measures” the measures which are absolutely continuous with respect to the Lebesgue measure and by $\mathcal{P}_c^{ac}(\mathbb{R}^d)$ the subset of $\mathcal{P}_c(\mathbb{R}^d)$ of AC measures.

Definition 2.1. For $\mu, \nu \in \mathcal{P}_c(\mathbb{R}^d)$, we denote by $\Pi(\mu, \nu)$ the set of *transference plans* from μ to ν , i.e. the probability measures on $\mathbb{R}^d \times \mathbb{R}^d$ with first marginal μ and second marginal ν . Let $p \in [1, \infty)$ and $\mu, \nu \in \mathcal{P}_c(\mathbb{R}^d)$. Define

$$W_p(\mu, \nu) := \inf_{\pi \in \Pi(\mu, \nu)} \left\{ \left(\int_{\mathbb{R}^d \times \mathbb{R}^d} \|x - y\|^p d\pi(x, y) \right)^{1/p} \right\}, \quad (11)$$

$$W_\infty(\mu, \nu) := \inf \{ \pi - \text{esssup} \|x - y\| : \pi \in \Pi(\mu, \nu) \}. \quad (12)$$

For $p \in [1, \infty]$, this quantity is called the **Wasserstein distance** or p -Wasserstein distance.

The idea behind this definition is the problem of *optimal transportation*, consisting in finding the optimal way to transport mass from a given measure to another. For a thorough introduction, see e.g. [49].

Between two microscopic configurations, we will use the following distance.

Definition 2.2. Let $p \in [1, \infty]$. Consider two configurations $X^0 := \{x_1^0, \dots, x_n^0\}$ and $X^1 := \{x_1^1, \dots, x_n^1\}$ of \mathbb{R}^d . Define the Wasserstein distance between X^0 and X^1 as follows:

$$W_p(X^0, X^1) := W_p(\mu^0, \mu^1),$$

where $\mu^0 := \frac{1}{n} \sum_i \delta_{x_i^0}$ and $\mu^1 := \frac{1}{n} \sum_i \delta_{x_i^1}$.

In this case, the Wasserstein distance can be rewritten as follows:

Proposition 1. Let $p \in [1, \infty)$. Consider two configurations $X^0 := \{x_1^0, \dots, x_n^0\}$ and $X^1 := \{x_1^1, \dots, x_n^1\}$ of \mathbb{R}^d . It holds

$$\begin{cases} W_p(X^0, X^1) = \inf_{\sigma \in \mathbb{S}_n} \left(\sum_{i=1}^n \frac{1}{n} \|x_i^0 - x_{\sigma(i)}^1\|^p \right)^{1/p}, \\ W_\infty(X^0, X^1) = \inf_{\sigma \in \mathbb{S}_n} \max_{i \in \{1, \dots, n\}} \|x_i^0 - x_{\sigma(i)}^1\|, \end{cases}$$

where \mathbb{S}_n is the set of permutations on $\{1, \dots, n\}$.

For a proof, we refer to [49, p. 5]. The idea behind this result is exactly the one of optimal transportation in the discrete setting: one has n initial and final configurations to be paired, with a given cost. The cost minimizer is the minimizer among all the permutations.

The Wasserstein distance satisfies some useful properties.

Proposition 2 (see [49, Chap. 7] and [20]). *It holds:*

1. For each pair $\mu, \nu \in \mathcal{P}_c(\mathbb{R}^d)$, the infimum in (11) or (12) is achieved.
2. For $p \in [1, \infty]$, the quantity W_p is a distance on $\mathcal{P}_c(\mathbb{R}^d)$. Moreover, for $p \in [1, \infty)$, the topology induced by the Wasserstein distance W_p on $\mathcal{P}_c(\mathbb{R}^d)$ coincides with the weak topology of measures, i.e. for all sequence $\{\mu_k\}_{k \in \mathbb{N}^*} \subset \mathcal{P}_c(\mathbb{R}^d)$ and all $\mu \in \mathcal{P}_c(\mathbb{R}^d)$, the following statements are equivalent:

$$(i) \quad W_p(\mu_k, \mu) \xrightarrow[k \rightarrow \infty]{} 0.$$

$$(ii) \quad \mu_k \xrightarrow[k \rightarrow \infty]{} \mu \text{ in the weak sense.}$$

The Wasserstein distance can be extended to all pairs of measures μ, ν compactly supported with the same total mass $\mu(\mathbb{R}^d) = \nu(\mathbb{R}^d) \neq 0$ by the formula

$$W_p(\mu, \nu) = |\mu|^{1/p} W_p \left(\frac{\mu}{|\mu|}, \frac{\nu}{|\nu|} \right).$$

In the rest of the paper, the following properties of the Wasserstein distance will be helpful.

Proposition 3 (see [49]). *Let $p \in [1, \infty]$ and μ, ν be two positive measures satisfying $\mu(\mathbb{R}^d) = \nu(\mathbb{R}^d)$ supported in a subset X . It then holds*

$$W_p(\mu, \nu) \leq \mu(\mathbb{R}^d) \text{diam}(X).$$

Proposition 4 (see [42, 49]). *Let μ, ρ, ν, η be four positive measures compactly supported satisfying $\mu(\mathbb{R}^d) = \nu(\mathbb{R}^d)$ and $\rho(\mathbb{R}^d) = \eta(\mathbb{R}^d)$.*

(i) *For each $p \in [1, \infty]$, it holds*

$$W_p^p(\mu + \rho, \nu + \eta) \leq W_p^p(\mu, \nu) + W_p^p(\rho, \eta).$$

(ii) *For each $p_1, p_2 \in [1, \infty]$ with $p_1 \leq p_2$, it holds*

$$\begin{cases} W_{p_1}(\mu, \nu) \leq W_{p_2}(\mu, \nu), \\ W_{p_2}(\mu, \nu) \leq \text{diam}(X)^{1-p_1/p_2} W_{p_1}^{p_1/p_2}(\mu, \nu), \end{cases} \quad (13)$$

where X contains the supports of μ and ν .

2.2 Well-posedness of System (2)

In this section, we study the macroscopic model (2), together with its connections with the Wasserstein distance. Consider the following system

$$\begin{cases} \partial_t \mu + \nabla \cdot (w\mu) = 0 & \text{in } \mathbb{R}^d \times \mathbb{R}^+, \\ \mu(\cdot, 0) = \mu^0 & \text{in } \mathbb{R}^d, \end{cases} \quad (14)$$

where $w : \mathbb{R}^d \times \mathbb{R}^+ \rightarrow \mathbb{R}^d$ is a time-dependent vector field. This equation is called the **continuity equation**. We now introduce the flow associated to System (14).

Definition 2.3. We define the **flow** associated to a vector field $w : \mathbb{R}^d \times \mathbb{R}^+ \rightarrow \mathbb{R}^d$ satisfying the Carathéodory condition as the application $(x^0, t) \mapsto \Phi_t^w(x^0)$ such that, for all $x^0 \in \mathbb{R}^d$, $t \mapsto \Phi_t^w(x^0)$ is the unique solution to

$$\begin{cases} \dot{x}(t) = w(x(t), t) \text{ for a.e. } t \geq 0, \\ x(0) = x^0. \end{cases} \quad (15)$$

It is classical that, for a vector field w satisfying the Carathéodory condition, System (15) is well-posed. See for instance [9].

We denote by Γ the set of the Borel maps $\gamma : \mathbb{R}^d \rightarrow \mathbb{R}^d$. We recall the definition of the *push-forward* of a measure.

Definition 2.4. For a $\gamma \in \Gamma$, we define the *push-forward* $\gamma\#\mu$ of a measure μ of \mathbb{R}^d as follows:

$$(\gamma\#\mu)(E) := \mu(\gamma^{-1}(E)),$$

for every subset E such that $\gamma^{-1}(E)$ is μ -measurable.

We now recall a standard result linking (14) and (15), known as the method of characteristics.

Theorem 2.5 (see [49, Th. 5.34]). *Let $T > 0$, $\mu^0 \in \mathcal{P}_c(\mathbb{R}^d)$ and w a vector field satisfying the Carathéodory condition. Then, System (14) admits a unique solution μ in $\mathcal{C}^0([0, T]; \mathcal{P}_c(\mathbb{R}^d))$, where $\mathcal{P}_c(\mathbb{R}^d)$ is equipped with the weak topology. Moreover:*

(i) *It holds $\mu(\cdot, t) = \Phi_t^w\#\mu^0$, where Φ_t^w is the flow of w as in Definition 2.3;*

(ii) If $\mu^0 \in \mathcal{P}_c^{ac}(\mathbb{R}^d)$, then $\mu(\cdot, t) \in \mathcal{P}_c^{ac}(\mathbb{R}^d)$.

We also recall the following proposition:

Proposition 5 (see [41, Prop. 4]). *Let $\mu, \nu \in \mathcal{P}_c(\mathbb{R}^d)$ and $w : \mathbb{R}^d \times \mathbb{R} \rightarrow \mathbb{R}^d$ be a vector field satisfying the Carathéodory condition, with a Lipschitz constant equal to L . For each $t \in \mathbb{R}$ and $p \in [1, \infty)$, it holds*

$$W_p(\Phi_t^w \# \mu, \Phi_t^w \# \nu) \leq e^{L|t|} W_p(\mu, \nu).$$

2.3 Approximate and exact controllability of System (1) and System (2)

In this section, we recall the main results of [24] about approximate and exact controllability of microscopic and macroscopic models. We first recall the precise notions of approximate and exact controllability.

Definition 2.6. We say that

- The microscopic system (1) (resp. the macroscopic system (2)) is **approximately controllable** from X^0 to X^1 (resp. from μ^0 to μ^1) at time T if for each $\varepsilon > 0$ there exists a sequence of controls $\{\mathbb{1}_\omega u_k\}_{k \in \mathbb{N}^*}$ satisfying the Carathéodory condition such that, denoting by X_k (resp. μ_k) the corresponding solution to System (1) (resp. System (2)), $X_k(T)$ converges to X^1 (resp. $\mu_k(T)$ converges weakly to μ^1).
- The microscopic system (1) (resp. the macroscopic system (2)) is **exactly controllable** from X^0 to X^1 (resp. from μ^0 to μ^1) at time T if there exists a control $\mathbb{1}_\omega u$ satisfying the Carathéodory condition such that the corresponding solution X (resp. μ) to System (1) (resp. System (2)) satisfies

$$X(T) = X^1 \text{ (resp. } \mu(T) = \mu^1 \text{)}.$$

For approximate controllability of System (2), the following result holds.

Theorem 2.7 (see [24]). *Let $\mu^0, \mu^1 \in \mathcal{P}_c^{ac}(\mathbb{R}^d)$ satisfying the Geometric Condition 1. Then there exists T such that System (2) is **approximately controllable** from μ^0 to μ^1 at time T with a control $\mathbb{1}_\omega u : \mathbb{R}^d \times \mathbb{R}^+ \rightarrow \mathbb{R}^d$ satisfying the Carathéodory condition.*

Remark 2. For exact controllability, the picture is completely different. The Carathéodory condition implies that the flow $\Phi_t^{v+\mathbb{1}_\omega u}$ is an homeomorphism. Since general μ^0, μ^1 are not homeomorphic, one needs to search a control with less regularity. The drawback is that one loses the uniqueness of the solution to System (2). This is the meaning of our recent result Theorem 2.8 below, proving that a class of controls ensuring exact controllability exists, but uniqueness is lost.

Theorem 2.8 (see [24]). *Let $\mu^0, \mu^1 \in \mathcal{P}_c^{ac}(\mathbb{R}^d)$ satisfying the Geometric Condition 1. Then, there exists $T > 0$ such that System (2) is **exactly controllable** from μ^0 to μ^1 at time T in the following sense: there exists a couple $(\mathbb{1}_\omega u, \mu)$ composed of a Borel vector field $\mathbb{1}_\omega u : \mathbb{R}^d \times \mathbb{R}^+ \rightarrow \mathbb{R}^d$ and a time-evolving measure μ being weak solution to System (2) and satisfying $\mu(T) = \mu^1$.*

Using Proposition 2, the approximate controllability of System (2) can be rewritten in terms of the Wasserstein distance:

Proposition 6. *The macroscopic system (2) is approximately controllable from μ^0 to μ^1 at time T if for each $\varepsilon > 0$ there exists a control $\mathbb{1}_\omega u$ satisfying the Carathéodory condition such that the corresponding solution μ to System (2) satisfies*

$$W_p(\mu^1, \mu(T)) \leq \varepsilon. \tag{16}$$

Remark 3. Properties (13) imply that all the Wasserstein distances W_p are equivalent for measures compactly supported and $p \in [1, \infty)$, see [49]. Thus, we can replace (16) by

$$W_1(\mu^1, \mu(T)) \leq \varepsilon.$$

Thus, in this work, we study approximate controllability by considering the distance W_1 only. We will use the distances W_2 and W_∞ , in some other specific cases only.

3 Infimum time for microscopic models

In this section, we prove Theorem 1.2 and 1.3, *i.e.* the infimum time for the approximate and exact controllability of microscopic models. We first obtain the following result:

Proposition 7. *Let $X^0 := \{x_1^0, \dots, x_n^0\} \subset \mathbb{R}^d$ and $X^1 := \{x_1^1, \dots, x_n^1\} \subset \mathbb{R}^d$ be two disjoint configurations (see Definition 1.1). Assume that ω is a bounded open connected set, the empirical measures associated to X^0 and X^1 (see (3)) satisfy the Geometric Condition 1 and the velocities v and $\mathbb{1}_\omega u$ satisfy the Carathéodory Condition 2. Consider the sequences $\{t_i^0\}_i$ and $\{t_i^1\}_i$ given in (5). Then*

$$\widetilde{M}_e(X^0, X^1) := \min_{\sigma \in \mathbb{S}_n} \max_{i \in \{1, \dots, n\}} |t_i^0 + t_{\sigma(i)}^1| \quad (17)$$

is the infimum time $T_e(X^0, X^1)$ to exactly control System (1) in the sense of Theorem 1.2.

Proof. We first prove the result corresponding to **Item (i)** of Theorem 1.2. Let $T := \widetilde{M}_e(X^0, X^1) + \delta$ with $\delta > 0$. Using the Geometric Condition 1, for all $i \in \{1, \dots, n\}$, there exist $s_i^0 \in (t_i^0, t_i^0 + \delta/3)$ and $s_i^1 \in (t_i^1, t_i^1 + \delta/3)$ such that

$$y_i^0 := \Phi_{s_i^0}^v(x_i^0) \in \omega \text{ and } y_i^1 := \Phi_{-s_i^1}^v(x_i^1) \in \omega.$$

This part of the proof is divided into two steps:

- In Step 1, we build a permutation σ and a flow on ω sending y_i^0 to $y_{\sigma(i)}^1$ for all $i \in \{1, \dots, n\}$ with no intersection of trajectories.
- In Step 2, we define a corresponding control sending x_i^0 to $x_{\sigma(i)}^1$ for all $i \in \{1, \dots, n\}$.

Item (i), Step 1: We first assume that ω is convex. The goal is to build a flow with no intersection of the characteristic. For all $i, j \in \{1, \dots, n\}$, we define the cost

$$K_{ij}(y_i^0, s_i^0, y_j^1, s_j^1) := \begin{cases} \|(y_i^0, s_i^0) - (y_j^1, T - s_j^1)\|_{\mathbb{R}^{d+1}} & \text{if } s_i^0 < T - s_j^1, \\ \infty & \text{otherwise.} \end{cases} \quad (18)$$

Consider the minimization problem:

$$\inf_{\pi \in \mathbb{B}_n} \frac{1}{n} \sum_{i,j=1}^n K_{ij}(y_i^0, s_i^0, y_j^1, s_j^1) \pi_{ij}, \quad (19)$$

where \mathbb{B}_n is the set of the bistochastic $n \times n$ matrices, *i.e.* the matrices $\pi := (\pi_{ij})_{1 \leq i, j \leq n}$ satisfying, for all $i, j \in \{1, \dots, n\}$,

$$\sum_{i=1}^n \pi_{ij} = 1, \quad \sum_{j=1}^n \pi_{ij} = 1, \quad \pi_{ij} \geq 0.$$

The infimum in (19) is finite since $T > \widetilde{M}_e(X^0, X^1)$. The problem (19) is a linear minimization problem on the closed convex set \mathbb{B}_n . Hence, as a consequence of Krein-Milman's Theorem (see [34]), the functional (19) admits a minimum at an extremal point of \mathbb{B}_n , *i.e.* a permutation matrix.

Let σ be a permutation, for which the associated matrix minimizes (19). Consider the straight trajectories $y_i(t)$ steering y_i^0 at time s_i^0 to $y_{\sigma(i)}^1$ at time $T - s_{\sigma(i)}^1$, that are explicitly defined by

$$y_i(t) := \frac{T - s_{\sigma(i)}^1 - t}{T - s_{\sigma(i)}^1 - s_i^0} y_i^0 + \frac{t - s_i^0}{T - s_{\sigma(i)}^1 - s_i^0} y_{\sigma(i)}^1. \quad (20)$$

We now prove by contradiction that these trajectories have no intersection: Assume that there exist i and j such that the associated trajectories $y_i(t)$ and $y_j(t)$ intersect. If we associate y_i^0 and y_j^0 to $y_{\sigma(j)}^0$ and $y_{\sigma(i)}^0$

respectively, *i.e.* we consider the permutation $\sigma \circ \mathcal{T}_{i,j}$, where $\mathcal{T}_{i,j}$ is the transposition between the i -th and the j -th elements, then the associated cost (19) is strictly smaller than the cost associated to σ . Indeed, using some geometric considerations in the space/time set (see Figure 3), we obtain

$$\left\{ \begin{array}{l} \|(y_i^0, s_i^0) - (y_{\sigma \circ \mathcal{T}_{i,j}(i)}^1, T - s_{\sigma \circ \mathcal{T}_{i,j}(i)}^1)\|_{\mathbb{R}^{d+1}} = \|(y_i^0, s_i^0) - (y_{\sigma(j)}^1, T - s_{\sigma(j)}^1)\|_{\mathbb{R}^{d+1}} \\ \qquad \qquad \qquad < \|(y_i^0, s_i^0) - (y_{\sigma(i)}^1, T - s_{\sigma(i)}^1)\|_{\mathbb{R}^{d+1}}, \\ \|(y_j^0, s_j^0) - (y_{\sigma \circ \mathcal{T}_{i,j}(j)}^1, T - s_{\sigma \circ \mathcal{T}_{i,j}(j)}^1)\|_{\mathbb{R}^{d+1}} = \|(y_j^0, s_j^0) - (y_{\sigma(i)}^1, T - s_{\sigma(i)}^1)\|_{\mathbb{R}^{d+1}} \\ \qquad \qquad \qquad < \|(y_j^0, s_j^0) - (y_{\sigma(j)}^1, T - s_{\sigma(j)}^1)\|_{\mathbb{R}^{d+1}}. \end{array} \right.$$

Here, we denote with $\|\cdot\|_{\mathbb{R}^{d+1}}$ the Euclidean norm in \mathbb{R}^{d+1} . This is in contradiction with the fact that σ




Figure 3: An optimal permutation.

minimizes (19).

Assume now that ω is not convex, but just open connected, hence arc-connected. Then, one replaces the norm $\|(y_i^0, s_i^0) - (y_j^1, T - s_j^1)\|_{\mathbb{R}^{d+1}}$ in (18) with the distance $d_{\bar{\omega}}(y_i^0, y_j^1) + |T - s_j^1 - s_i^0|$, where

$$d_{\bar{\omega}}(y_i^0, y_j^1) = \inf_{\gamma \in C^1([0,1]; \mathbb{R}^d)} \left\{ \int_0^1 |\dot{\gamma}(t)| dt \quad \text{s.t.} \quad \gamma(0) = y_i^0, \quad \gamma(1) = y_j^1, \quad \gamma(t) \in \bar{\omega} \quad \forall t \in [0,1] \right\}.$$

For each pair, infimum is finite and it is realized by some curve γ_{ij} , since $\bar{\omega}$ is compact and arc-connected. For the global problem (18), there exist a minimizing permutation σ steering y_i^0 at time s_i^0 to $y_{\sigma(i)}^1$ at time $T - s_{\sigma(i)}^1$. The corresponding curves $\gamma_{i\sigma(i)}$ are not intersecting, for the same reasoning explained above for convex sets. Then, since ω is open, one can deform them to find paths in ω that are arbitrarily close to the $\gamma_{i\sigma(i)}$, still not intersecting and C^∞ functions of time. We denote by $y_i(t)$ such deformed path steering y_i^0 at time s_i^0 to $y_{\sigma(i)}^1$ at time $T - s_{\sigma(i)}^1$.

Item (i), Step 2: Consider now the following trajectories z_i :

$$z_i(t) := \begin{cases} \Phi_t^v(x_i^0) & \text{for all } t \in (0, s_i^0), \\ y_i(t) & \text{for all } t \in (s_i^0, T - s_{\sigma(i)}^1), \\ \Phi_{t-T}^v(x_i^1) & \text{for all } t \in (T - s_{\sigma(i)}^1, T), \end{cases}$$

where $y_i(t)$ are defined in Item (i), Step 1. The trajectories z_i have no intersection. By construction of trajectories $y_i(t)$, it holds $z_i(t) \in \omega$ for all $t \in (s_i^0, T - s_{\sigma(i)}^1)$. For all $i \in \{1, \dots, n\}$, choose r_i, R_i satisfying $0 < r_i < R_i$ and such that it holds

$$\begin{cases} B_{r_i}(z_i(t)) \subset B_{R_i}(z_i(t)) \subset \omega & \text{for all } t \in (s_i^0, T - s_{\sigma(i)}^1) \\ B_{R_i}(z_i(t)) \cap B_{R_j}(z_j(t)) = \emptyset & \text{for all } t \in (0, T), \end{cases}$$

with $i, j \in \{1, \dots, n\}$. Such radii r_i, R_i exist as a consequence of the fact that we deal with a finite number of trajectories that do not cross. The corresponding control can be chosen as a function that is piecewise

constant with respect to the time variable and C^∞ with respect to the x variable given by:

$$u(x, t) := \begin{cases} \dot{y}_i(t) - v & \text{if } t \in (s_i^0, T - s_{\sigma(i)}^1) \text{ and } x \in B_{r_i}(z_i(t)), \\ 0 & \text{if } t \in (s_i^0, T - s_{\sigma(i)}^1) \text{ and } x \notin B_{R_i}(z_i(t)), \\ C^\infty\text{-space-spline} & \text{if } t \in (s_i^0, T - s_{\sigma(i)}^1) \text{ and } x \in B_{R_i}(z_i(t)) \setminus B_{r_i}(z_i(t)), \\ 0 & \text{if } t \notin (s_i^0, T - s_{\sigma(i)}^1), \end{cases}$$

We have built a control satisfying the Carathéodory condition such that each i -th component of the associated solution to System (1) is $z_i(t)$, hence steering x_i^0 to $x_{\sigma(i)}^1$ in time T .

We now prove the result corresponding to **Item (ii)** of Theorem 1.2. Assume that System (1) is exactly controllable at a time $T > M_e^*(X^0, X^1)$, and consider σ the corresponding permutation defined by $x_i(T) = x_{\sigma(i)}^1$. The idea of the proof is that the trajectory steers x_i^0 to ω in time t_i^0 , then it moves inside ω for a small but positive time, then it steers a point from ω to $x_{\sigma(i)}^1$ in time $t_{\sigma(i)}^1$, hence $T > t_i^0 + t_{\sigma(i)}^1$.

Fix an index $i \in \{1, \dots, n\}$. First recall the definition of $t_i^0, t_{\sigma(i)}^1$ and observe that it holds both $T > t_i^0$ and $T > t_{\sigma(i)}^1$. Then, the trajectory $x_i(t)$ satisfies¹ $x_i(t) \notin \omega$ for all $t \in (0, t_i^0)$, as well as $x_i(t) \notin \omega$ for all $t \in (T - t_{\sigma(i)}^1, T)$. Moreover, we prove that it exists $\tau_i \in (0, T)$ for which it holds $x_i(\tau_i) \in \omega$. By contradiction, if such τ_i does not exist, then the trajectory $x_i(t)$ never crosses the control region, hence it coincides with $\Phi_t^v(x_i^0)$. But in this case, by definition of t_i^0 as the infimum of times such that $\Phi_t^v(x_i^0) \in \omega$ and recalling that $t_i^0 < T$, there exists $\tau_i \in (t_i^0, T)$ such that it holds $x_i(\tau_i) = \Phi_{\tau_i}^v(x_i^0) \in \omega$. Contradiction. Also observe that ω is open, hence there exists ϵ_i such that $x_i(\tau) \in \omega$ for all $\tau \in (\tau_i - \epsilon_i, \tau_i + \epsilon_i)$. We merge the conditions $x_i(t) \notin \omega$ for all $t \in (0, t_i^0) \cup (T - t_{\sigma(i)}^1, T)$ with $x_i(\tau) \in \omega$ for all $\tau \in (\tau_i - \epsilon_i, \tau_i + \epsilon_i)$ with a given $\tau_i \in (0, T)$. This implies that it holds $t_i^0 < \tau_i < T - t_{\sigma(i)}^1$, hence

$$T > t_i^0 + t_{\sigma(i)}^1.$$

Such estimate holds for any $i \in \{1, \dots, n\}$. Using the definition of $\widetilde{M}_e(X^0, X^1)$, we deduce that $T > \widetilde{M}_e(X^0, X^1)$.

We finally prove the result corresponding to **Item (iii)** of Theorem 1.2. By definition of $M_e^*(X^0, X^1)$, there exists $l \in \{0, 1\}$ and $m \in \{1, \dots, n\}$ such that $M_e^*(X^0, X^1) = t_m^l$. We only study the case $l = 0$, since the case $l = 1$ can be recovered by reversing time. We consider the trajectory starting from such x_m^0 only. By definition of t_m^0 , the trajectory $\Phi_t^v(x_m^0)$ satisfies $\Phi_t^v(x_m^0) \notin \omega$ for all $t \in [0, M_e^*(X^0, X^1)]$. Then, for any choice of the control u localized in ω , it holds $\Phi_t^{v+1\omega u}(x_m^0) = \Phi_t^v(x_m^0)$, *i.e.* the choice of the control plays no role in the trajectory starting from x_m^0 on the time interval $t \in [0, M_e^*(X^0, X^1)]$. Observe that it holds $v(\Phi_t^v(x_m^0)) \neq 0$ for all $t \in [0, M_e^*(X^0, X^1)]$, due to the fact that the vector field is time-independent and the trajectory $\Phi_t^v(x_m^0)$ enters ω for some $t > M_e^*(X^0, X^1)$.

We now prove that the set of times $t \in [0, M_e^*(X^0, X^1)]$ for which exact controllability holds is finite. A necessary condition to have exact controllability at time t is that the equation $\Phi_t^v(x_m^0) = x_i^1$ admits a solution for some time $t \in [0, M_e^*(X^0, X^1)]$ and index $i \in \{1, \dots, n\}$. Then, we aim to prove that the set of times-indexes (t, i) solving such equation is finite. By contradiction, assume to have an infinite number of solutions (t, i) . Since the set $i \in \{1, \dots, n\}$ is finite, this implies that there exists an index I and an infinite number of (distinct) times $t_k \in [0, M_e^*(X^0, X^1)]$ such that $\Phi_{t_k}^v(x_m^0) = x_I^1$. Since v is an autonomous vector field, this implies that, for each pair t_{k_1}, t_{k_2} it holds $\Phi_{t_{k_1}-t_{k_2}}^v(x_I^1) = x_I^1$, hence $\Phi_t^v(x_I^1)$ is a periodic trajectory, with period $|t_{k_1} - t_{k_2}|$. By compactness of $[0, M_e^*(X^0, X^1)]$, there exists a converging subsequence (that we do not relabel) $t_k \rightarrow t_* \in [0, M_e^*(X^0, X^1)]$. Then $\Phi_{t_*}^v(x_I^1)$ is a periodic trajectory with arbitrarily small period, hence x_I^1 is an equilibrium². This contradicts the fact that for all $t \in [0, M_e^*(X^0, X^1)]$ it holds $v(\Phi_t^v(x_m^0)) \neq 0$. \square

¹These estimates hold even in the specific case of $x_i^0 \in \omega$, for which it holds $t_i^0 = 0$.

²Such final statement can be easily proved by contradiction: if $v(x_I^1) \neq 0$, then apply the local rectifiability theorem for Lipschitz vector field [11] and prove that any eventually periodic trajectory passing through x_I^1 has a strictly positive minimal period.

Remark 4. Proposition 7 can be interpreted as follows: Each particle at point x_i^0 needs to be sent on a target point $x_{\sigma(i)}^1$. The time $\widetilde{M}_e(X^0, X^1)$ coincides with the infimum time for a particle at x_i^0 to enter in ω and then go from ω to $x_{\sigma(i)}^1$. We are thus assuming that the particle travels with an arbitrarily large speed in ω .

Formula (17) leads to the proof of Theorem 1.2.

Proof of Theorem 1.2. Let σ_0 be a minimizing permutation in (17). We build recursively a sequence of permutations $\{\sigma_1, \dots, \sigma_n\}$ as follows:

- Let k_1 be such that $t_{\sigma_0(k_1)}^1$ is a maximum of $\{t_{\sigma_0(1)}^1, \dots, t_{\sigma_0(n)}^1\}$. We denote by $\sigma_1 := \sigma_0 \circ \mathcal{T}_{1, k_1}$, where, for all $i, j \in \{1, \dots, n\}$, $\mathcal{T}_{i, j}$ is the transposition between the i -th and the j -th elements. It holds

$$\left\{ \begin{array}{l} t_{k_1}^0 + t_{\sigma_0(k_1)}^1 \geq t_1^0 + t_{\sigma_0(1)}^1, \\ t_{k_1}^0 + t_{\sigma_0(k_1)}^1 \geq t_1^0 + t_{\sigma_0(k_1)}^1 = t_1^0 + t_{\sigma_0 \circ \mathcal{T}_{1, k_1}(1)}^1 = t_1^0 + t_{\sigma_1(1)}^1, \\ t_{k_1}^0 + t_{\sigma_0(k_1)}^1 \geq t_{k_1}^0 + t_{\sigma_0(1)}^1 = t_{k_1}^0 + t_{\sigma_0 \circ \mathcal{T}_{1, k_1}(k_1)}^1 = t_{k_1}^0 + t_{\sigma_1(k_1)}^1. \end{array} \right.$$

Thus σ_1 minimizes (17) too, since it holds

$$\max_{i \in \{1, \dots, n\}} \{t_i^0 + t_{\sigma_0(i)}^1\} \geq \max_{i \in \{1, \dots, n\}} \{t_i^0 + t_{\sigma_1(i)}^1\}.$$

- Assume that σ_j is built. Let k_{j+1} be such that $t_{\sigma_j(k_{j+1})}^1$ is a maximum of $\{t_{\sigma_j(j+1)}^1, \dots, t_{\sigma_j(n)}^1\}$. Again, we clearly have

$$\left\{ \begin{array}{l} t_{k_{j+1}}^0 + t_{\sigma_j(k_{j+1})}^1 \geq t_{j+1}^0 + t_{\sigma_j(j+1)}^1, \\ t_{k_{j+1}}^0 + t_{\sigma_j(k_{j+1})}^1 \geq t_{j+1}^0 + t_{\sigma_j(k_{j+1})}^1 = t_{j+1}^0 + t_{\sigma_j \circ \mathcal{T}_{j+1, k_{j+1}}(j+1)}^1 \\ \quad = t_{j+1}^0 + t_{\sigma_{j+1}(j+1)}^1, \\ t_{k_{j+1}}^0 + t_{\sigma_j(k_{j+1})}^1 \geq t_{k_{j+1}}^0 + t_{\sigma_j(k_j)}^1 = t_{k_{j+1}}^0 + t_{\sigma_j \circ \mathcal{T}_{j+1, k_{j+1}}(k_{j+1})}^1 \\ \quad = t_{k_{j+1}}^0 + t_{\sigma_{j+1}(k_{j+1})}^1. \end{array} \right.$$

Thus $\sigma_{j+1} := \sigma_j \circ \mathcal{T}_{j+1, k_{j+1}}$ minimizes (17) too:

$$\max_{i \in \{1, \dots, n\}} \{t_i^0 + t_{\sigma_j(i)}^1\} \geq \max_{i \in \{1, \dots, n\}} \{t_i^0 + t_{\sigma_{j+1}(i)}^1\}.$$

The sequence $\{t_{\sigma_n(1)}^1, \dots, t_{\sigma_n(n)}^1\}$ is then decreasing and σ_n is a minimizing permutation in (17). We deduce that $\widetilde{M}_e(X^0, X^1) = M_e(X^0, X^1)$. \square

With Theorem 1.2, we give an explicit and simple expression of the infimum time for microscopic models. This result is useful in numerical simulations of Section 5 (in particular, see Algorithm 2). Let μ^0 and μ^1 be the probability measures defined by

$$\mu^0 := \sum_{i=1}^n \frac{1}{n} \delta_{x_i^0} \text{ and } \mu^1 := \sum_{i=1}^n \frac{1}{n} \delta_{x_i^1}, \quad (21)$$

where the points x_i^0 (resp. x_i^1) are disjoint. We now deduce that $T_e(X^0, X^1)$ is equal to $S(\mu^0, \mu^1)$ given in (8), when μ^0 and μ^1 are given by (21).

Corollary 1. *Let $X^0 := \{x_1^0, \dots, x_n^0\} \subset \mathbb{R}^d$ and $X^1 := \{x_1^1, \dots, x_n^1\} \subset \mathbb{R}^d$ be two disjoint configurations. Consider μ^0 and μ^1 the corresponding measures given in (21). Assume that ω is a bounded open connected set, μ^0, μ^1 satisfy the Geometric Condition 1 and the velocities v and $\mathbb{1}_\omega u$ satisfy the Carathéodory Condition 2.*

Then the infimum time $T_e(X^0, X^1)$ is equal to $S(\mu^0, \mu^1)$ given in (8).

Proof. Remark that if the sequences $\{t_i^0\}_{i \in \{1, \dots, n\}}$ and $\{t_i^1\}_{i \in \{1, \dots, n\}}$ are non-decreasingly and non-increasingly ordered respectively, then for all $m \in (\frac{i-1}{n}, \frac{i}{n}]$ it holds

$$\mathcal{F}_{\mu^0}^{-1}(m) = t_i^0 \quad \text{and} \quad \mathcal{B}_{\mu^1}^{-1}(1-m) = t_i^1.$$

Then, the result is given by identification of the expression of $M_e(X^0, X^1)$ given in (6) with the expression of $S(\mu^0, \mu^1)$ in (8). \square

We now prove Theorem 1.3, which characterizes the infimum time for approximate control of System (1).

Proof of Theorem 1.3. We first prove **Item (i)**. Consider $X^0 := \{x_1^0, \dots, x_n^0\}$ and $X^1 := \{x_1^1, \dots, x_n^1\}$ two disjoint configurations satisfying the Geometric Condition 1. We first prove that the infimum time $T_a(X^0, X^1)$ is equal to

$$\widetilde{M}_a(X^0, X^1) := \min_{\sigma \in \mathbb{S}_n} \max_{i \in \{1, \dots, n\}} |t_i^0 + \bar{t}_{\sigma(i)}^1|.$$

First assume that $T > \widetilde{M}_a(X^0, X^1)$. Let $\varepsilon > 0$. For each x_i^1 , we need to find points y_i^1 satisfying

$$\|y_i^1 - x_i^1\| \leq \varepsilon \quad \text{and} \quad y_i := \Phi_{-\bar{t}_i^1}^v(y_i^1) \in \omega. \quad (22)$$

For each x_i^1 , observe that the Geometric Condition 1 implies that either $x_i^1 \in \omega$ or that the trajectory enters ω backward in time. In the first case, define $y_i^1 := x_i^1$. In the second case, remark that $v(\Phi_{-t}^v(x_i^1))$ is nonzero for a whole interval $t \in [0, \bar{t}]$, with $\bar{t} > \bar{t}_i^1$, and $\Phi_{-\bar{t}_i^1}^v(x_i^1) \in \bar{\omega}$, hence the flow $\Phi_{-\bar{t}_i^1}^v(\cdot)$ is an homeomorphism in a neighborhood of x_i^1 . Then, there exists $y_i^1 \in \mathbb{R}^d$ such that (22) is satisfied.

We denote by $Y^1 := \{y_1^1, \dots, y_n^1\}$. For all $i \in \{1, \dots, n\}$, since $y_i \in \omega$, then $t^1(y_i^1) \leq \bar{t}_i^1$, hence $\widetilde{M}_e(X^0, Y^1) \leq \widetilde{M}_a(X^0, X^1) < T$. Proposition 7 implies that we can exactly steer X^0 to Y^1 at time T with a control u satisfying the Carathéodory condition. Denote by $X(t)$ the solution to System (1) for the initial condition X^0 and the control u . It holds

$$W_1(X^1, X(T)) = W_1(X^1, Y^1) \leq \sum_{i=1}^n \frac{1}{n} \|y_i^1 - x_i^1\| \leq \varepsilon. \quad (23)$$

Choose now $\varepsilon = \frac{1}{k}$ for each $k \in \mathbb{N}^*$, and denote by u_k the corresponding control and with $x_{i,k}(t)$ the associated solution to System (1) for the i -th particle. By the definition of W_1 in the discrete case and (23), there exists a permutation $\sigma_k \in \mathbb{S}_n$ for which it holds

$$\|x_{i,k}(T) - x_{\sigma_k(i),k}^1\| \leq \frac{n}{k},$$

for all $i = 1, \dots, n$. Since the space of permutations \mathbb{S}_n is finite, one can extract a subsequence with σ_k constant. With such subsequence, the statement (i) is proved.

We now prove **Item (ii)**. Consider a time $T \geq M_a^*(X^0, X^1)$ at which System (1) is approximately controllable. We aim to prove that it satisfies $T > \widetilde{M}_a(X^0, X^1)$. For each $k \in \mathbb{N}^*$, there exists a control u_k satisfying the Carathéodory condition such that the corresponding solution $X_k(t)$ to System (2) satisfies

$$W_1(X^1, X_k(T)) \leq \frac{1}{k}. \quad (24)$$

We denote by $Y_k^1 := \{y_{k,1}^1, \dots, y_{k,n}^1\}$ the configuration defined by

$$y_{k,i}^1 := X_{k,i}(T),$$

where $X_{k,i}$ is the i -th component of X_k . Since X^0 is disjoint and u_k satisfies the Carathéodory condition, then Y_k^1 is disjoint too. We now prove that it holds

$$T > M_e^*(X^0, Y_k^1). \quad (25)$$

Since $T > M_a^*(X^0, X^1)$, then (25) is equivalent to $T > t_i^1(y_{k,i}^1)$ for all $i \in \{1, \dots, n\}$. By contradiction, assume that there exists $j \in \{1, \dots, n\}$ such that

$$t^1(y_{k,j}^1) \geq T.$$

We distinguish two cases:

- If $t^1(y_{k,j}^1) > T$, then for any $t \in [0, T]$ it holds $\Phi_{-t}^v(y_{k,j}^1) \notin \omega$. Thus, the localized control does not act on the trajectory, *i.e.* for each $t \in [0, T]$ it holds $\Phi_{-t}^v(y_{k,j}^1) = \Phi_{-t}^{v+\mathbb{1}_\omega u_k}(y_{k,j}^1)$. Since $y_{k,j}^1 = \Phi_T^{v+\mathbb{1}_\omega u_k}(x_j^0) = \Phi_T^v(x_j^0)$, then $\Phi_t^v(x_j^0) \notin \omega$ for all $t \in [0, T]$. This is a contradiction with the fact that $t_j^0 \leq M_a^*(X^0, X^1) < T$. Thus (25) holds.
- If $t^1(y_{k,j}^1) = T$, then for all $t \in [0, T)$ it holds $\Phi_{-t}^v(y_{k,j}^1) \notin \omega$. Since ω is open, then it also holds $\Phi_{-T}^v(y_{k,j}^1) \notin \omega$. We then conclude as in the previous case.

Since $Y_k^1 = X_k(T)$, then Proposition 7 implies that

$$T > \widetilde{M}_e(X^0, Y_k^1). \quad (26)$$

For each control u_k , denote by σ_k the permutation for which $y_{k,i}^1 = \Phi_T^{v+\mathbb{1}_\omega u_k}(x_{\sigma_k(i)}^0)$. Up to extract a subsequence, for all k large enough, σ_k is equal to a permutation σ . Inequality (24) implies that for all $i \in \{1, \dots, n\}$ it holds

$$y_{k,i}^1 \xrightarrow[k \rightarrow \infty]{} x_{\sigma(i)}^1. \quad (27)$$

Since $t^1(y_{k,i}^1) \leq \widetilde{M}_e(X^0, Y_k^1) < T$, up to a subsequence, for a $s_i \geq 0$, it holds

$$t^1(y_{k,i}^1) \xrightarrow[k \rightarrow \infty]{} s_i. \quad (28)$$

Using (27), (28) and the continuity of the flow, it holds

$$\begin{aligned} & \|\Phi_{-t^1(y_{k,i}^1)}^v(y_{k,i}^1) - \Phi_{-s_i}^v(x_{\sigma(i)}^1)\| \\ & \leq \|\Phi_{-t^1(y_{k,i}^1)}^v(y_{k,i}^1) - \Phi_{-s_i}^v(y_{k,i}^1)\| + \|\Phi_{-s_i}^v(y_{k,i}^1) - \Phi_{-s_i}^v(x_{\sigma(i)}^1)\| \xrightarrow[k \rightarrow \infty]{} 0. \end{aligned}$$

The fact that $\Phi_{-t^1(y_{k,i}^1)}^v(y_{k,i}^1) \in \bar{\omega}$ for each $i = 1, \dots, n$ leads to $\Phi_{-s_i}^v(x_{\sigma(i)}^1) \in \bar{\omega}$. Thus

$$\bar{t}^1(x_{\sigma(i)}^1) \leq \lim_{k \rightarrow \infty} t^1(y_{k,i}^1).$$

Denoting by $\delta := (T - \widetilde{M}_e(X^0, X^1))/2$ and using (26), there exists $K \in \mathbb{N}$ such that for all $k > K$ it holds

$$\begin{aligned} \widetilde{M}_a(X^0, X^1) &= \min_{\bar{\sigma} \in \mathbb{S}_n} \max_{i \in \{1, \dots, n\}} |t_i^0 + \bar{t}_{\bar{\sigma}(i)}^1| \\ &\leq \max_{i \in \{1, \dots, n\}} |t_i^0 + \bar{t}_{\sigma(i)}^1| \\ &\leq \max_{i \in \{1, \dots, n\}} |t_i^0 + t^1(y_{k,\sigma(i)}^1)| + \delta \\ &= \widetilde{M}_e(X^0, Y_k^1) + \delta < T. \end{aligned}$$

We finally prove **Item (iii)** of Theorem 1.3. Let $T \in (0, M_a^*(X^0, X^1))$ be such that System (1) is approximately controllable. For any $\varepsilon > 0$, there exists u_ε such that the associated trajectory to System (1) satisfies

$$W_1(X_\varepsilon(T), X^1) < \varepsilon. \quad (29)$$

There exists $j \in \{1, \dots, n\}$ s.t. $t^0(x_j^0) = M_a^*(X^0, X^1) > T$ or $\bar{t}^1(x_j^1) = M_a^*(X^0, X^1) > T$. We distinguish these two cases:

- Assume that $t^0(x_j^0) = M_a^*(X^0, X^1) > T$.

Define $x_{\varepsilon,j}(t) := \Phi_t^{v+1\omega u_\varepsilon}(x_j^0)$. Inequality (29) implies that there exists $k \in \{1, \dots, n\}$ such that

$$\|x_{\varepsilon,j}(T) - x_{k_\varepsilon}^1\| < \varepsilon. \quad (30)$$

As $t^0(x_j^0) > T$, the trajectory $\Phi_t^v(x_j^0)$ does not cross the control set ω for $t \in [0, T]$, hence $x_{\varepsilon,j}(T) = \Phi_T^{v+1\omega u_\varepsilon}(x_j^0) = \Phi_T^v(x_j^0)$ does not depend on ε . Define $R := \frac{1}{2} \min_{p,q} \|x_p^1 - x_q^1\|$, that is strictly positive since X^1 is disjoint. For each $\varepsilon < R$, Estimate (30) gives $k_\varepsilon = k$ independent on ε and

$$x_{\varepsilon,j}(T) = \Phi_t^v(x_j^0) = x_k^1.$$

Use now the proof of Item (iii) in Proposition 7 to prove that the equation $\Phi_t^v(x_j^0) = x_k^1$ admits a finite number of solutions (t, k) with $t \in [0, t^0(x_j^0)]$ and $k \in \{1, \dots, n\}$.

- Assume now that $\bar{t}^1(x_j^1) = M_a^*(X^0, X^1) > T$. Again, inequality (29) implies that there exists $k_\varepsilon \in \{1, \dots, n\}$ such that

$$\|x_{\varepsilon,k_\varepsilon}(T) - x_j^1\| = \|\Phi_T^{v+1\omega u_\varepsilon}(x_{k_\varepsilon}^0) - x_j^1\| < \varepsilon.$$

As $\bar{t}^1(x_j^1) > T$, the trajectory $\Phi_{-t}^v(x_j^1)$ does not cross the control set $\bar{\omega}$ for all $t \in [0, T]$. Then for all $t \in [0, T]$, there exists $r(t) > 0$ such that $B_{r(t)}(\Phi_{-t}^v(x_j^1)) \cap \omega = \emptyset$. By compactness of the set $\{\Phi_{-t}^v(x_j^1) : t \in [0, T]\}$, there exist $t_1, \dots, t_N \in [0, T]$ for which it holds

$$\{\Phi_{-t}^v(x_j^1) : t \in [0, T]\} \subset \bigcup_{i=1}^N B_{r(t_i)}(\Phi_{-t_i}^v(x_j^1)) \subset \omega^c.$$

Thus there exists a common $r > 0$ such that $B_r(\Phi_{-t}^v(x_j^1)) \cap \omega = \emptyset$ for all $t \in [0, T]$. For $\varepsilon < r$, it holds $k_\varepsilon = k$ and

$$x_{\varepsilon,k}(T) = \Phi_T^v(x_k^0) = x_j^1.$$

We conclude as in the previous case.

Finally, apply the permutation method used in the proof of Theorem 1.2 to prove that it holds $\widetilde{M}_a(X^0, X^1) = M_a(X^0, X^1)$. \square

Remark 5. We illustrate Item (iii) of Theorem 1.2 by giving an example in which System (1) is never exactly controllable on $[0, M_e^*(X^0, X^1))$ and another where System (1) is exactly controllable at a time $T \in [0, M_e^*(X^0, X^1))$.

Figure 4 (left): Consider $\omega := (-1, 1) \times (-1.5, 1.5)$, $v := (1, 0)$, $X^0 := (-2, 0)$ and $X^1 := (2, 0)$. The time $M_e^*(X^0, X^1)$ at which we can act on the particles and the minimal time $M_e(X^0, X^1)$ are respectively equal to 1 and 2. We observe that System (1) is neither exactly controllable nor approximately controllable on the interval $[0, M_e^*(X^0, X^1))$.

Figure 4 (right): Consider $\omega := (-2, 0) \times (-1.5, 1.5)$, the vector field $v(x, y) = (-y, x)$ and X^0, X^1 as follows

$$X^0 := \{(\sqrt{2}/2, -\sqrt{2}/2)\}, \quad X^1 := \{(\sqrt{2}/2, \sqrt{2}/2)\}.$$

The time $M_e^*(X^0, X^1)$ at which we can act on the particles and the minimal time $M_e(X^0, X^1)$ are respectively equal to $3\pi/4$ and π . We remark that System (1) is exactly controllable, then approximately controllable at time $T = \pi/2 \in [0, M_e^*(X^0, X^1))$.




Figure 4: Examples in the case $T \in (0, M_e^*(X^0, X^1))$.

4 Infimum time for macroscopic models

This section is devoted to the proof of main Theorem 1.4 about infimum time for AC measures. We first introduce the notion of infimum time up to small mass in Section 4.1. We then give some comparisons between the microscopic and macroscopic cases in Section 4.2. We finally use the obtained results to prove Theorem 1.4 in Section 4.3.

4.1 Infimum time in the microscopic setting up to small masses

In this section, we introduce the notion of infimum time up to small mass and prove some results about this notion in the microscopic case.

From now on, we denote by $\lfloor x \rfloor$ the floor of $x \in \mathbb{R}$.

Definition 4.1 (Infimum time up to small mass). Consider μ^0, μ^1 be two AC-measures compactly supported with the same total mass $\mu^0(\mathbb{R}^d) = \mu^1(\mathbb{R}^d) = \gamma > 0$. Recall that $S^*(\mu^0, \mu^1)$ is defined by (7). We denote by $T_{e,\varepsilon}(\mu^0, \mu^1)$ the infimum time to exactly steer μ^0 to μ^1 up to a mass ε . More precisely, it is the infimum of times $T \geq S^*(\mu^0, \mu^1)$ for which there exists a control $\mathbb{1}_{\omega} u : \mathbb{R}^d \times \mathbb{R}^+ \rightarrow \mathbb{R}^d$ satisfying the Carathéodory condition and two sets R^0, R^1 of \mathbb{R}^d such that $\mu^0(R^0) = \mu^1(R^1) = \varepsilon$ and

$$\Phi_T^{v+\mathbb{1}_{\omega} u} \# \mu_{|(R^0)^c}^0 = \mu_{|(R^1)^c}^1.$$

We similarly define the minimal time $T_{a,\varepsilon}(\mu^0, \mu^1)$ to approximately steer μ^0 to μ^1 up to a mass ε .

Let $\gamma > 0$ and $X^0 := \{x_1^0, \dots, x_n^0\} \subset \mathbb{R}^d$, $X^1 := \{x_1^1, \dots, x_n^1\} \subset \mathbb{R}^d$ be two disjoint configurations (see Definition 1.1). We denote by $T_{e,\varepsilon}(X^0, X^1, \gamma)$ the infimum time to exactly steer X^0 to X^1 up to a mass ε for a total mass γ . More precisely, it is the infimum of times $T \geq M_e^*(X^0, X^1)$ such that there exists a control $\mathbb{1}_{\omega} u : \mathbb{R}^d \times \mathbb{R}^+ \rightarrow \mathbb{R}^d$ satisfying the Carathéodory condition and two permutations $\sigma_0, \sigma_1 \in \mathbb{S}_n$ with $R := \lfloor \frac{n\varepsilon}{\gamma} \rfloor$ such that

$$x_{\sigma_0(i)}(T) = x_{\sigma_1(i)},$$

for all $i \in \{1, \dots, n - R\}$. We similarly define $T_{a,\varepsilon}(X^0, X^1, \gamma)$ the minimal time to approximately steer X^0 to X^1 up to a mass ε for a total mass γ .

Remark 6. If $\varepsilon < \gamma/n$, then the infimum time up to a small mass clearly coincides with the standard infimum time studied in Section 3.

Remark 7. Let $\gamma > 0$ and $X^0 := \{x_1^0, \dots, x_n^0\} \subset \mathbb{R}^d$, $X^1 := \{x_1^1, \dots, x_n^1\} \subset \mathbb{R}^d$ be two disjoint configurations (see Definition 1.1). We remark that

$$T_{e,\varepsilon}(X^0, X^1, \gamma) = T_{e,\varepsilon}(\mu^0, \mu^1)$$

for the measures μ^0 and μ^1 given by

$$\mu^0 := \sum_{i=1}^n \frac{\gamma}{n} \delta_{x_i^0} \text{ and } \mu^1 := \sum_{i=1}^n \frac{\gamma}{n} \delta_{x_i^1}. \quad (31)$$

For all pair of measures μ^0 and μ^1 compactly supported with the same total mass $\gamma := \mu^0(\mathbb{R}^d) = \mu^1(\mathbb{R}^d)$, we define the quantity

$$S_\varepsilon(\mu^0, \mu^1) := \sup_{m \in [0, \gamma - \varepsilon]} \{\mathcal{F}_{\mu^0}^{-1}(m) + \mathcal{B}_{\mu^1}^{-1}(\gamma - \varepsilon - m)\}. \quad (32)$$

We have the following result:

Proposition 8. *Let $\gamma > 0$ and $X^0 := \{x_1^0, \dots, x_n^0\}, X^1 := \{x_1^1, \dots, x_n^1\} \subset \mathbb{R}^d$ be two disjoint configurations. Assume that ω is a bounded open connected set, the empirical measures associated to X^0 and X^1 (see (3)) satisfy the Geometric Condition 1 and the velocities v and $\mathbb{1}_\omega u$ satisfy the Carathéodory Condition 2. Assume that the sequences $\{t_i^0\}_{i \in \{1, \dots, n\}}$ and $\{t_i^1\}_{i \in \{1, \dots, n\}}$ are respectively non-decreasingly and non-increasingly ordered. The infimum time $T_{e, \varepsilon}(X^0, X^1, \gamma)$ to exactly steer μ^0 to μ^1 up to a mass ε for a total mass γ is equal to*

$$M_{e, \varepsilon}(X^0, X^1, \gamma) := \max_{1 \leq i \leq n-R} \{t^0(x_i^0) + t^1(x_{i+R}^1)\},$$

where $R := \lfloor n\varepsilon\gamma \rfloor$. Moreover, it is also equal to

$$S_\varepsilon(X^0, X^1, \gamma) := S_\varepsilon(\mu^0, \mu^1), \quad (33)$$

where the measures μ^0 and μ^1 are defined in (31).

Proof. We can adapt the proof of Proposition 7 as follows. We first replace the set \mathbb{B}_n of bistochastic matrices in (19) by the set $\mathbb{B}_{n, R}$ composed with the matrices satisfying

$$\sum_i \pi_{ij} \leq 1, \quad \sum_j \pi_{ij} \leq 1, \quad \sum_{ij} \pi_{ij} = n - R \text{ and } \pi_{ij} \geq 0.$$

The set $\mathbb{B}_{n, R}$ is closed and convex. Consider the minimization problem

$$\inf_{\pi \in \mathbb{B}_{n, R}} \frac{1}{n} \sum_{i, j=1}^n K_{ij}(y_i^0, s_i^0, y_j^1, s_j^1) \pi_{ij}, \quad (34)$$

where the quantities $K_{ij}(y_i^0, s_i^0, y_j^1, s_j^1)$ are defined in (18). Also in this case, the infimum in (34) is finite since $T > \widetilde{M}_{e, \varepsilon}(X^0, X^1, \gamma)$. Problem (34) is linear, hence, again as a consequence of Krein-Milman's Theorem (see [34]), some minimisers of this functional are extremal points, that are matrices composed of a permutation sub-matrix for some rows and columns, and zeros for other rows and columns. We then define

$$\widetilde{M}_{e, \varepsilon}(X^0, X^1, \gamma) := \min_{\sigma_0, \sigma_1 \in \mathbb{S}_n} \max_{1 \leq i \leq n-R} |t_{\sigma_0(i)}^0 + t_{\sigma_1(i)}^1|.$$

By applying the permutation method of the proof of Theorem 1.2, we prove that $M_{e, \varepsilon}(X^0, X^1, \gamma)$ is equal to $\widetilde{M}_{e, \varepsilon}(X^0, X^1, \gamma)$.

Following the lines of Corollary 1, this also implies that formula (33) holds, by using the definitions of $\mathcal{F}_{\mu^0}^{-1}$ and $\mathcal{B}_{\mu^1}^{-1}$. \square

4.2 Comparison of microscopic and macroscopic cases

The goal of this section is to give some estimates of the functions S^* and S_ε defined in (7) and (32), depending both on their arguments μ^0, μ^1 and the control region. They will be instrumental to prove Theorem 1.4 in Section 4.3. For this reason, we will specify the considered control region in the notation $t^0, t^1, \mathcal{F}_{\mu^0}, \mathcal{B}_{\mu^1}, \mathcal{F}_{\mu^0, \omega}^{-1}$ and $\mathcal{B}_{\mu^1, \omega}^{-1}$ in an index, i.e. $t_\omega^0, t_\omega^1, \mathcal{F}_{\mu^0, \omega}, \mathcal{B}_{\mu^1, \omega}, \mathcal{F}_{\mu^0, \omega}^{-1}$ and $\mathcal{B}_{\mu^1, \omega}^{-1}$. For the functions S, S_ε and S^* , the control region will be specified as follows: $S(\mu^0, \mu^1, \omega), S_\varepsilon(\mu^0, \mu^1, \omega)$ and $S^*(\mu^0, \mu^1, \omega)$.

Proposition 9. Let $\mu^0, \mu^1 \in \mathcal{P}_c^{ac}(\mathbb{R}^d)$. Assume that ω is a bounded open connected set, μ^0, μ^1 satisfy the Geometric Condition 1 and the velocities v and $\mathbb{1}_\omega u$ satisfy the Carathéodory Condition 2. Fix $\{f_n\}_{n \in \mathbb{N}}$ be a decreasing sequence of positive numbers converging to zero. For each $n \in \mathbb{N}$, let ω_n be defined by

$$\omega_n := \{x \in \mathbb{R}^d : d(x, \omega^c) > f_n\}. \quad (35)$$

Let two sequences $\{\mu_n^0\}_{n \in \mathbb{N}}, \{\mu_n^1\}_{n \in \mathbb{N}}$ of compactly supported measures be given, satisfying the Geometric Condition 1 with respect to ω . Consider also two sequences of Borel sets $\{R_n^0\}_{n \in \mathbb{N}}, \{R_n^1\}_{n \in \mathbb{N}}$ of \mathbb{R}^d that satisfy

$$\begin{cases} r_n := \mu^0(R_n^0) = \mu^1(R_n^1) \xrightarrow{n \rightarrow \infty} 0, \\ \mu_n^0(\mathbb{R}^d) = \mu_n^1(\mathbb{R}^d) = 1 - r_n, \\ W_\infty(\mu|_{(R_n^0)^c}, \mu_n^0) < f_n, \quad W_\infty(\mu|_{(R_n^1)^c}, \mu_n^1) < f_n. \end{cases} \quad (36)$$

Let $S_\varepsilon(\cdot, \cdot, \tilde{\omega})$ be the function defined in (32) for a given control set $\tilde{\omega}$. Then for each $\varepsilon, \delta > 0$, there exists $N \in \mathbb{N}^*$ such that for all $n \geq N$, it holds

$$(i) \quad S_{2\varepsilon}(\mu_n^0, \mu_n^1, \omega_n) \leq S_\varepsilon(\mu^0, \mu^1, \omega) + \delta.$$

$$(ii) \quad S_{2\varepsilon}(\mu^0, \mu^1, \omega) \leq S_\varepsilon(\mu_n^0, \mu_n^1, \omega_n) + \delta.$$

Moreover, there exist two sequences of Borel sets $U_{0,n}, U_{1,n} \subset \mathbb{R}^d$ such that

$$\begin{cases} \mu_n^0(U_{0,n}) = \mu_n^1(U_{1,n}) \xrightarrow{n \rightarrow \infty} 1, \\ \limsup_{n \rightarrow \infty} S^*(\mu_n^0|_{U_{0,n}}, \mu_n^1|_{U_{1,n}}, \omega_n) \leq S^*(\mu^0, \mu^1, \omega). \end{cases}$$

Proof. We first prove **Item (i)**. Fix $\varepsilon, \delta > 0$. For each $n \in \mathbb{N}$, define

$$V_n^0 := \{x^0 \in \text{supp}(\mu^0) : \Phi_t^v(B_{f_n}(x^0)) \subset \subset \omega_n \text{ for a } t \in [0, t_\omega^0(x^0) + \delta/2]\}.$$

We define

$$r_n^0 := \mu^0((V_n^0)^c). \quad (37)$$

Since $\{f_n\}_{n \in \mathbb{N}^*}$ is decreasing, then $\{\omega_n\}_{n \in \mathbb{N}^*}$ is a sequence of increasing sets, hence V_n^0 is a sequence of increasing sets. Thus r_n^0 is decreasing, hence $\lim_{n \rightarrow \infty} r_n^0$ exists. We now prove that it holds

$$\lim_{n \rightarrow \infty} r_n^0 = 0. \quad (38)$$

We prove it by contradiction. Assume that $\lim_{n \rightarrow \infty} r_n^0 > 0$. Then it holds $\mu^0((V_n^0)^c) > C > 0$ for all $n \in \mathbb{N}^*$. In particular, the limit set $(\bigcap_{n=1}^\infty (V_n^0)^c) \cap \text{supp}(\mu^0)$ is non-empty, hence there exists $x^0 \in \text{supp}(\mu^0)$ such that

$$\Phi_t^v(B_{f_n}(x^0)) \not\subset \omega_n, \quad (39)$$

for all $t \in [0, t_\omega^0(x^0) + \delta/2]$ and $n \in \mathbb{N}^*$. Due to the Geometric Condition 1, such time $t_\omega^0(x^0)$ is finite. Since ω is open, for a $t_\omega^*(x^0) \in (t_\omega^0(x^0), t_\omega^0(x^0) + \delta/2)$ and a $r(x^0) > 0$, it holds $B_{r(x^0)}(\Phi_{t_\omega^*}^v(x^0)) \subset \subset \omega$. By continuity of Φ_t^v , there exists $\hat{r}(x^0) > 0$ such that

$$\Phi_{t_\omega^*}^v(B_{\hat{r}(x^0)}(x^0)) \subset \subset \omega. \quad (40)$$

Since (39) and (40) are in contradiction, for n large enough, we conclude that (38) holds. We deduce that, for all $x^0 \in V_n^0$, it holds

$$\begin{aligned} \xi^0 \in \overline{B_{f_n}(x^0)} &\Rightarrow \Phi_t^v(\xi^0) \in \omega_n \text{ for some } t \in [0, t_\omega^0(x^0) + \delta/2] \\ &\Rightarrow t_{\omega_n}^0(\xi^0) \leq t_\omega^0(x^0) + \delta/2. \end{aligned} \quad (41)$$

For each $n \in \mathbb{N}^*$, consider an optimal transference plan π_n realizing the ∞ -Wasserstein distance $W_\infty(\mu_{|(R_n^0)^c}^0, \mu_n^0)$, see Proposition 2. We remark that (36) implies that

$$|x^0 - \xi^0| < f_n \text{ for } \pi_n\text{-almost every } (x^0, \xi^0) \in (R_n^0)^c \cap \text{supp}(\mu^0) \times \text{supp}(\mu_n^0). \quad (42)$$

Thus, combining (41) and (42), it holds

$$t_{\omega_n}^0(\xi^0) \leq t_\omega^0(x^0) + \delta/2 \quad (43)$$

for π_n -almost every pair (x^0, ξ^0) with $x^0 \in (R_n^0)^c \cap V_n^0$ and $\xi^0 \in \text{supp}(\mu_n^0)$. Using (37) and (43), we obtain

$$\begin{aligned} \mathcal{F}_{\mu^0, \omega}(t) &\leq \mu^0(\{x^0 \in (R_n^0)^c \cap V_n^0 : t_\omega^0(x^0) \leq t\}) + \mu^0(R_n^0) + \mu^0((V_n^0)^c) \\ &= \pi_n(\{(x^0, \xi^0) \in ((R_n^0)^c \cap V_n^0) \times \text{supp}(\mu_n^0) : t_\omega^0(x^0) \leq t\}) + r_n + r_n^0 \\ &\leq \pi_n(\{(x^0, \xi^0) \in ((R_n^0)^c \cap V_n^0) \times \text{supp}(\mu_n^0) : t_{\omega_n}^0(\xi^0) \leq t + \delta/2\}) + r_n + r_n^0 \\ &\leq \mu_n^0(\{\xi^0 \in \text{supp}(\mu_n^0) : t_{\omega_n}^0(\xi^0) \leq t + \delta/2\}) + r_n + r_n^0 \\ &= \mathcal{F}_{\mu_n^0, \omega_n}(t + \delta/2) + r_n + r_n^0, \end{aligned}$$

for all $t \in \mathbb{R}^+$. Similarly, we have

$$\mathcal{B}_{\mu^1, \omega}(t) \leq \mathcal{B}_{\mu_n^1, \omega_n}(t + \delta/2) + r_n + r_n^1,$$

for all $t \in \mathbb{R}^+$, where r_n^1 is defined similarly to (37). We deduce that the generalized inverse satisfies

$$\begin{aligned} \mathcal{F}_{\mu_n^0, \omega_n}^{-1}(m) &:= \inf\{t \geq 0 : \mathcal{F}_{\mu_n^0, \omega_n}(t) \geq m\} \leq \inf\{t \geq \delta/2 : \mathcal{F}_{\mu^0, \omega}(t - \delta/2) - r_n^0 - r_n \geq m\} \\ &= \inf\{s \geq 0 : \mathcal{F}_{\mu^0, \omega}(s) \geq m + r_n^0 + r_n\} + \delta/2 = \mathcal{F}_{\mu^0, \omega}^{-1}(m + r_n^0 + r_n) + \delta/2, \end{aligned}$$

for all $m \in (0, 1 - r_n^0 - r_n)$. Similarly, we obtain

$$\mathcal{B}_{\mu_n^1, \omega_n}^{-1}(1 - r_n - 2\varepsilon - m) \leq \mathcal{B}_{\mu^1, \omega}^{-1}(1 + r_n^1 - 2\varepsilon - m) + \delta/2,$$

for all $m \in (r_n^1 - 2\varepsilon, 1 - r_n - 2\varepsilon)$. For n large enough, we have

$$\begin{aligned} S_{2\varepsilon}(\mu_n^0, \mu_n^1, \omega_n) &:= \sup_{m \in [0, 1 - r_n - 2\varepsilon]} \{\mathcal{F}_{\mu_n^0, \omega_n}^{-1}(m) + \mathcal{B}_{\mu_n^1, \omega_n}^{-1}(1 - r_n - 2\varepsilon - m)\} \\ &\leq \sup_{m \in [0, 1 - r_n - 2\varepsilon]} \{\mathcal{F}_{\mu^0, \omega}^{-1}(m + r_n^0 + r_n) + \mathcal{B}_{\mu^1, \omega}^{-1}(1 + r_n^1 - 2\varepsilon - m)\} + \delta \\ &\leq \sup_{m \in [r_n^0 + r_n, 1 + r_n^0 - 2\varepsilon]} \{\mathcal{F}_{\mu^0, \omega}^{-1}(m) + \mathcal{B}_{\mu^1, \omega}^{-1}(1 + r_n^1 + r_n^0 + r_n - 2\varepsilon - m)\} + \delta. \end{aligned}$$

Observe that it holds $0 \leq r_n^0 + r_n$. For n large enough, it also holds $1 + r_n^0 - 2\varepsilon \leq 1 - \varepsilon$, as well as $r_n^1 + r_n^0 + r_n \leq \varepsilon$. Since $\mathcal{B}_{\mu^1, \omega}^{-1}$ is increasing, this implies

$$S_{2\varepsilon}(\mu_n^0, \mu_n^1, \omega_n) \leq \sup_{m \in [0, 1 - \varepsilon]} \{\mathcal{F}_{\mu^0, \omega}^{-1}(m) + \mathcal{B}_{\mu^1, \omega}^{-1}(1 - \varepsilon - m)\} + \delta = S_\varepsilon(\mu^0, \mu^1, \omega) + \delta$$

for n sufficiently large.

We now prove **Item (ii)**. For all $n \in \mathbb{N}$, define

$$\tilde{V}_n^0 := \{x^0 \in \text{supp}(\mu_n^0) : \Phi_t^v(B_{f_n}(x^0)) \subset\subset \omega \text{ for a } t \leq t_{\omega_n}^0(x^0) + \delta/2\}$$

and $\tilde{r}_n^0 := \mu^0((\tilde{V}_n^0)^c)$. Using the same argument as for item (i), we can prove that

$$t_\omega^0(\xi^0) \leq t_{\omega_n}^0(x^0) + \delta/2 \quad (44)$$

for π_n -almost every $x^0 \in \tilde{V}_n^0$ and $\xi^0 \in \text{supp}(\mu^0) \cap (R_n^0)^c$, where π_n is a transportation plan realizing $W_\infty(\mu_n^0, \mu^0)$. Inequality (44) implies

$$\begin{aligned}
\mathcal{F}_{\mu_n^0, \omega_n}(t) &\leq \mu_n^0(\{x^0 \in \tilde{V}_n^0 : t_{\omega_n}^0(x^0) \leq t\}) + \mu^0((\tilde{V}_n^0)^c) \\
&= \pi_n(\{(x^0, \xi^0) \in \tilde{V}_n^0 \times (\text{supp}(\mu^0) \cap (R_n^0)^c) : t_{\omega_n}^0(x^0) \leq t\}) + \tilde{r}_n^0 \\
&\leq \pi_n(\{(x^0, \xi^0) \in \tilde{V}_n^0 \times (\text{supp}(\mu^0) \cap (R_n^0)^c) : t_{\omega}^0(\xi^0) \leq t + \delta/2\}) + \tilde{r}_n^0 \\
&\leq \mu^0(\{\xi^0 \in \text{supp}(\mu^0) \cap (R_n^0)^c : t_{\omega}^0(\xi^0) \leq t + \delta/2\}) + \tilde{r}_n^0 \\
&\leq \mu^0(\{\xi^0 \in \text{supp}(\mu^0) : t_{\omega}^0(\xi^0) \leq t + \delta/2\}) + \mu^0(R_n^0) + \tilde{r}_n^0 \\
&= \mathcal{F}_{\mu^0, \omega}(t + \delta/2) + r_n + \tilde{r}_n^0,
\end{aligned}$$

for all $t \in \mathbb{R}^+$. We also have

$$\mathcal{B}_{\mu_n^1, \omega_n}(t) \leq \mathcal{B}_{\mu^1, \omega}(t + \delta/2) + r_n + \tilde{r}_n^1,$$

for all $t \in \mathbb{R}^+$, where \tilde{r}_n^1 is similarly defined. We conclude as before.

The last statement holds for $U_{0,n} := V_n^0$ and $U_{1,n} := V_n^1$, where

$$V_n^1 := \{x^1 \in \text{supp}(\mu^1) : \Phi_{-t}^v(B_{f_n}(x^1)) \subset \subset \omega_n \text{ for a } t \leq t_{\omega}^1(x^1) + \delta/2\}.$$

□

4.3 Proof of Theorem 1.4

In this section, we prove Theorem 1.4. The proof is based on the results obtained in Section 4.1 and 4.2. We highlight that the idea of the proof is to approximate the macroscopic crowds with microscopic ones with a sufficiently large number of agents, find a control steering one microscopic crowd to another, then adapt such control for the macroscopic one. For this reason, this result can be seen as the limit of Proposition 8.

Proof of Theorem 1.4. We first prove **Item (i)**. Fix $s > 0$ and $\varepsilon > 0$ and define the time

$$T := S(\mu^0, \mu^1, \omega) + s.$$

To prove the theorem, we aim to show that there exists an admissible control u such that

$$W_1(\mu^1, \Phi_T^{v+\mathbb{1}_{\omega}u} \# \mu^0) \leq \varepsilon.$$

We assume that the space dimension is $d := 2$, but the reader will see that the proof can be clearly adapted to any space dimension. The proof is divided into three steps.

Step 1: In this step, we discretize the measures, in four sub-steps. We first discretize uniformly in space the supports of μ^0 and μ^1 . To simplify the presentation, assume that $\text{supp}(\mu^0) \subset (0, 1)^2$ and $\text{supp}(\mu^1) \subset (0, 1)^2$. For all $k := (k_1, k_2) \in \{0, \dots, n-1\}^2$, consider the set $C_{n,k}$ defined by

$$C_{n,k} := \left[\frac{k_1}{n}, \frac{k_1+1}{n} \right] \times \left[\frac{k_2}{n}, \frac{k_2+1}{n} \right].$$

Then define $\mu_{nk}^l := \mu_{C_{n,k}}^l$ for $l = 0, 1$. Remark that the number of indexes nk is exactly n^2 .

We now consider only the decomposition of μ^0 , while the decomposition of μ^1 is similar. To send a measure to another, these measures need to have the same total mass. With this goal, we perform the second step of our decomposition: on each set $C_{n,k}$ consider a decomposition

$$a_0 = \frac{k_1}{n} < a_1 < \dots < a_m \leq \frac{k_1+1}{n}$$

such that it holds $\mu_{nk}^0([a_i, a_{i+1}] \times [\frac{k_2}{n}, \frac{k_2+1}{n}]) = \frac{1}{n^3}$ for all $i = 0, \dots, m-1$, as well as $\mu_{nk}^0([a_m, \frac{k_1+1}{n}] \times [\frac{k_2}{n}, \frac{k_2+1}{n}]) < \frac{1}{n^3}$. If $\mu^0(C_{n,k}) < \frac{1}{n^3}$, we set $m = 0$. One can always perform such a decomposition, since μ^0 is absolutely continuous. Then define $C_{nki}^0 := [a_i, a_{i+1}] \times [\frac{k_2}{n}, \frac{k_2+1}{n}]$ for $i = 0, \dots, m-1$ as well as $\mu_{nki}^0 := (\mu_{nk}^0)|_{C_{nki}^0}$. Also define

$$\bar{\mu}_{nk}^0 := (\mu_{nk}^0)|_{[a_m, \frac{k_1+1}{n}] \times [\frac{k_2}{n}, \frac{k_2+1}{n}]}.$$

The meaning of such decomposition is the following: each μ_{nki}^0 has mass $\frac{1}{n^3}$ and is localized in C_{nki}^0 , while $\bar{\mu}_{nk}^0$ has a mass strictly smaller than $\frac{1}{n^3}$ and is localized in $C_{n,k}^0$. Remark that the number of indexes i depends on n, k . Nevertheless, the total mass not contained in the sets C_{nki}^0 is smaller than $\frac{1}{n}$, hence the total number of indexes nki is between $n^3 - n^2$ and n^3 .

We now consider the third step of the decomposition: each μ_{nki}^0 is localized in the set $C_{nki}^0 := [a_i, a_{i+1}] \times [\frac{k_2}{n}, \frac{k_2+1}{n}]$. For each index nki , define a decomposition

$$a_{i0} = \frac{k_2}{n} < a_{i1} < \dots < a_{im} \leq \frac{k_2+1}{n}$$

such that it holds $\mu_{nki}^0([a_i, a_{i+1}] \times [a_{ij}, a_{i(j+1)}]) = \frac{1}{n^4}$ for each $j = 0, \dots, m-1$. Then define $C_{nki}^0 := [a_i, a_{i+1}] \times [a_{ij}, a_{i(j+1)}]$ for each $j = 0, \dots, m-1$, as well as $\mu_{nki}^0 := (\mu_{nki}^0)|_{C_{nki}^0}$. Remark that the number of indexes j depends on nki , but that the total number of indexes nki is between $n^4 - n^3$ and n^4 .

If the dimension of the space is larger than 2, one simply needs to decompose the μ_{nki}^0 with respect to the subsequent dimension with blocks of mass $\frac{1}{n^5}$, and so on, up to cover all dimensions.

The second and third step of the decomposition are represented in Figure 5.

For more details on such discretization, we refer to [24, Prop. 3.1].




Figure 5: Example of a partition of $C_{n,k}$ in cells such as C_{nki}^0 (hashed).

We finally consider the fourth step of our decomposition: in each cell $C_{nki}^0 = [a_i, a_{i+1}] \times [a_{ij}, a_{i(j+1)}]$, it is defined an absolutely continuous measure μ_{nki}^0 with mass $\frac{1}{n^4}$. Thus, it exists a constant $\epsilon > 0$ such that the set

$$B_{nki}^0 := [a_i + \epsilon, a_{i+1} - \epsilon] \times [a_{ij} + \epsilon, a_{i(j+1)} - \epsilon] \quad (45)$$

satisfies $\mu_{nki}^0(B_{nki}^0) = \frac{1}{n^4} - \frac{1}{n^6}$. See Figure 6 for a graphical description of this decomposition. For more details of such construction, we also refer to [24, Prop. 3.1].

We finally define $\tilde{\mu}_{nki}^0 := (\mu_{nki}^0)|_{B_{nki}^0}$ and $\bar{\mu}_{nki}^0 := (\mu_{nki}^0)|_{C_{nki}^0 \setminus B_{nki}^0}$.

We discretize similarly the measure μ^1 on sets $C_{nki}^1, C_{nki}^1, B_{nki}^1$, defining $\bar{\mu}_{nk}^1, \bar{\mu}_{nki}^1, \tilde{\mu}_{nki}^1$.

Step 2: In this step, we prove exact controllability of microscopic approximations $\hat{\mu}_n^0, \hat{\mu}_n^1$ of μ^0, μ^1 . In a first step, we define such sequence of approximations $\hat{\mu}_n^0, \hat{\mu}_n^1$ of μ^0, μ^1 satisfying hypotheses of Proposition 9.




Figure 6: Example of cells B_{nkij}^0 (hashed).

Step 2.1: We aim to define a sequence of microscopic approximations $\hat{\mu}_n^0, \hat{\mu}_n^1$ of μ^0, μ^1 , that satisfy hypotheses of Proposition 9. We first observe the following: the number of $nkij$ indexes are between $n^4 - n^3$ and n^4 both for μ^0 and μ^1 . We then choose exactly a set I_n^0 of $n^4 - n^3$ indexes $nkij$ for μ^0 , as well as a set I_n^1 of $n^4 - n^3$ indexes $nkij$ for μ^1 .

We then define the following measures:

$$\begin{aligned} \tilde{\mu}_n^0 &:= \sum_{nkij \in I_n^0} \tilde{\mu}_{nkij}^0, & \tilde{\mu}_n^1 &:= \sum_{nkij \in I_n^1} \tilde{\mu}_{nkij}^1, \\ \bar{\mu}_n^0 &:= \mu^0 - \tilde{\mu}_n^0 = \sum_{nk} \bar{\mu}_{nk}^0 + \sum_{nki} \bar{\mu}_{nki}^0 + \sum_{nkij} \bar{\mu}_{nkij}^0 + \sum_{nkij \notin I_n^0} \tilde{\mu}_{nkij}^0, \end{aligned} \quad (46)$$

and similarly $\tilde{\mu}_n^1, \bar{\mu}_n^1 = \mu^1 - \tilde{\mu}_n^1$. The idea of the notation is the following: the measure μ^0 is decomposed into a “relevant” part $\tilde{\mu}_n^0$ and a negligible one $\bar{\mu}_n^0$. Such decomposition only depends on the parameter n of the grid, as well as the choice of $n^4 - n^3$ indexes defining the set I_n^0 . The same holds for the decomposition of μ^1 .

We are ready to define the microscopic approximation $\hat{\mu}_n^0$. Observe that each $\tilde{\mu}_{nkij}^0$ is supported in the set B_{nkij}^0 defined in (45) and has mass $\frac{1}{n^4} - \frac{1}{n^6}$. For each $nkij \in I_n^0$, choose a point x_{nkij}^0 belonging to the support of μ_{nkij}^0 . Then define

$$\hat{\mu}_n^0 := \sum_{nkij \in I_n^0} \left(\frac{1}{n^4} - \frac{1}{n^6} \right) \delta_{x_{nkij}^0}.$$

Repeat the same construction for μ^1 , then defining $\hat{\mu}_n^1$.

Step 2.2: We now prove that sequences $\hat{\mu}_n^0, \hat{\mu}_n^1$ defined above, together with μ^0, μ^1 , satisfy the hypotheses of Proposition 9. Define the sequence

$$f_n := e^{LT} \sqrt{2}/n,$$

where L is the Lipschitz constant of the vector field v . Then observe that $x_{nkij}^0 \in \text{supp}(\mu^0)$ for all $nkij$, hence the fact that μ^0 satisfies the Geometric Condition 1 implies that $\hat{\mu}_n^0$ satisfies it too.

For each n , define the set R_n^0 as the support of $\bar{\mu}_n^0$. This implies that

$$r_n := \mu^0(R_n^0) = \bar{\mu}_n^0(\mathbb{R}^d) = 1 - \tilde{\mu}_n^0(\mathbb{R}^d) = 1 - (n^4 - n^3) \left(\frac{1}{n^4} - \frac{1}{n^6} \right) \rightarrow 0. \quad (47)$$

Moreover, by construction it holds $\hat{\mu}_n^0(\mathbb{R}^d) = \tilde{\mu}_n^0(\mathbb{R}^d) = 1 - r_n$. Finally, observe that each set B_{nkij}^0 has diameter strictly smaller than the diameter of C_{nkij}^0 , hence smaller than $\frac{\sqrt{2}}{n} < f_n$. Moreover, it is clear that it holds $(\mu^0)_{|(R_n^0)^c} = \tilde{\mu}_n^0$ by construction. Then, one can estimate $W_\infty((\mu^0)_{|(R_n^0)^c}, \hat{\mu}_n^0)$ by decomposing

³For the space \mathbb{R}^d , it is clearly sufficient to replace $\sqrt{2}$ with \sqrt{d} .

them in each set B_{nkij}^0 , *i.e.*

$$\begin{aligned} W_\infty((\mu^0)|_{(R_n^0)^c}, \hat{\mu}_n^0) &\leq \max_{nkij \in I_n^0} W_\infty(\tilde{\mu}_{nkij}^0, \hat{\mu}_{nkij}^0) \leq \\ &\max_{nkij \in I_n^0} \text{diam}(\text{supp}(\tilde{\mu}_{nkij}^0) \cup \text{supp}(\hat{\mu}_{nkij}^0)) \leq \max_{nkij \in I_n^0} \text{diam}(B_{nkij}^0) < f_n. \end{aligned}$$

Repeat the same procedure for μ^1 , defining R_n^1 and proving the same estimates with f_n, r_n . Then, all conditions of (36) are satisfied. Summing up, all hypotheses of Proposition 9 are satisfied.

Since v is uniformly bounded, there exists $R > 0$ independent on n and ε , such that

$$\bigcup_{t \in [0, T]} \Phi_t^v(\omega) \subset\subset B_R(0). \quad (48)$$

We now estimate the value of $S_{\varepsilon/4R}(\hat{\mu}_n^0, \hat{\mu}_n^1, \omega_n)$, where ω_n is defined in (35). Observe that it clearly holds

$$S_{\varepsilon/8R}(\mu^0, \mu^1, \omega) \leq S(\mu^0, \mu^1, \omega).$$

Apply Proposition 9 for $\delta := s/2$, that gives the following estimate:

$$S_{\varepsilon/4R}(\hat{\mu}_n^0, \hat{\mu}_n^1, \omega_n) + \frac{s}{2} \leq S_{\varepsilon/8R}(\mu^0, \mu^1, \omega) + s \leq S(\mu^0, \mu^1, \omega) + s = T,$$

for n large enough.

Step 2.3: We now prove existence of a control exactly steering $\hat{\mu}_n^0$ to $\hat{\mu}_n^1$ up to a mass $\varepsilon/4R$. Using Condition (9), ω_n is connected for n large enough, Using Proposition 8, it holds

$$T_{e, \varepsilon/4R}(\hat{\mu}_n^0, \hat{\mu}_n^1, \omega_n) = S_{\varepsilon/4R}(\hat{\mu}_n^0, \hat{\mu}_n^1, \omega_n) < T.$$

Then there exists $I_{n, \varepsilon}^l \subset I_n^0$ ($l = 0, 1$) such that

$$|I_{n, \varepsilon}^l| = (1 - \varepsilon/4R)(n^4 - n^3) \quad (49)$$

(assumed integer for simplicity), a bijection among indexes $\sigma_n : I_{n, \varepsilon}^0 \rightarrow I_{n, \varepsilon}^1$ and a control $\mathbb{1}_{\omega_n} \hat{u}_n$ satisfying the Carathéodory condition such that

$$\Phi_T^{v + \mathbb{1}_{\omega_n} \hat{u}_n}(x_{nkij}^0) = x_{\sigma_n(nkij)}^1$$

for all $nkij \in I_{n, \varepsilon}^0$. We define

$$\tilde{\mu}_{n, \varepsilon}^l := \sum_{nkij \in I_{n, \varepsilon}^l} \tilde{\mu}_{nkij}^l, \quad \bar{\mu}_{n, \varepsilon}^l := \mu^l - \tilde{\mu}_{n, \varepsilon}^l, \quad \hat{\mu}_{n, \varepsilon}^l := \sum_{nkij \in I_{n, \varepsilon}^l} \left(\frac{1}{n^4} - \frac{1}{n^6} \right) \delta_{x_{nkij}^l}.$$

For simplicity of notation, for each n we assume to re-arrange indexes $nkij$ both for $\hat{\mu}_{n, \varepsilon}^0$ and $\hat{\mu}_{n, \varepsilon}^1$, so that $I_{n, \varepsilon}^0 = I_{n, \varepsilon}^1$ and the bijection σ_n is the identity. We will then write from now on

$$\Phi_T^{v + \mathbb{1}_{\omega_n} \hat{u}_n}(x_{nkij}^0) = x_{nkij}^1. \quad (50)$$

We also denote by $\hat{x}_{nkij}(t)$ the corresponding trajectory, steering x_{nkij}^0 to x_{nkij}^1 in time T . Finally, we denote

$$t_{nkij}^0 = t^0(x_{nkij}^0, \omega_n), \quad t_{nkij}^1 = t^1(x_{nkij}^1, \omega_n).$$

We also make the following observation: the trajectory steering x_{nkij}^0 to x_{nkij}^1 is known to enter ω_n at time t_{nkij}^0 and to exit it at time $T - t_{nkij}^1$. Due to the construction of the control in the proof of Proposition 7, all trajectories $\hat{x}_{nkij}(t)$ are contained in ω_n for all times $t \in (t_{nkij}^0, T - t_{nkij}^1)$.

Step 3: We now prove approximate controllability from μ^0 to μ^1 in time $T = S(\mu^0, \mu^1, \omega) + s$. The idea is to write a control approximately steering μ^0 to μ^1 , based on the controls \hat{u}_n defined in (50) exactly steering $\hat{\mu}_{n, \varepsilon}^0$ to $\hat{\mu}_{n, \varepsilon}^1$ in time T .

Fix a given n . For each $nkij \in I_{n, \varepsilon}^0$ apply the following procedure:

1. the set B_{nkij}^0 evolves via the uncontrolled flow Φ_t^v in the time interval $[0, t_{nkij}^0]$; at this final time, the evolved set $B_{nkij}(t_{nkij}^0) := \Phi_{t_{nkij}^0}^v(B_{nkij}^0)$ is completely contained in ω .
2. we then apply a control w_{nkij} concentrating the mass around $\hat{x}_{nkij}(t)$ defined in (50) for the time interval $[t_{nkij}^0, T - t_{nkij}^1]$. Such mass is then moved to the corresponding evolved set $B_{nkij}(T - t_{nkij}^1)$ around $\hat{x}(T - t_{nkij}^1)$;
3. we finally let such set evolve via the uncontrolled flow Φ_t^v up to time T ; the set $B_{nkij}(T)$ is centered around $\hat{x}(T) = x_{nkij}^1$.

In the following, we will prove that such strategy provides the desired result. In the first step, we will precisely define the strategy and prove estimates about the evolution of sets. In the second and final step, we will provide estimates about the Wasserstein distance, to prove approximate controllability.

Step 3.1: We now define a control approximately steering μ^0 to μ^1 . We first define the control and the desired evolution for each index $nkij \in I_{n,\varepsilon}^0$. We recall that the control \hat{u}_n given in (50) defines the trajectory $\hat{x}_{nkij}(t)$ steering x_{nkij}^0 to x_{nkij}^1 . We then define an adapted control $\tilde{u}_n(t, x)$ that concentrates mass around such trajectory. We choose

$$\tilde{u}_{nkij}(t, x) = \hat{u}_n(t, \hat{x}_{nkij}(t)) + C_n(\hat{x}_{nkij}(t) - x),$$

where C_n is a positive constant that will be chosen later. It is clearly a linear feedback stabilizing the system around the trajectory \hat{x}_{nkij} . We also define a corresponding vector field

$$w_{nkij}(t, x) := v(\hat{x}_{nkij}(t)) + \tilde{u}_{nkij}(t, x). \quad (51)$$

Observe that in this formula the vector field v is evaluated at $\hat{x}_{nkij}(t)$ only, and not at the point x . We then define the evolution of the corresponding evolved cell $B_{nkij}(t)$ as follows:

$$B_{nkij}(t) = \begin{cases} \Phi_t^v(B_{nkij}^0) & \text{if } t \in [0, t_{nkij}^0], \\ \Phi_t^{w_{nkij}}(B_{nkij}(t_{nkij}^0)) & \text{if } t \in [t_{nkij}^0, T - t_{nkij}^1], \\ \Phi_t^v(B_{nkij}(T - t_{nkij}^1)) & \text{if } t \in [T - t_{nkij}^1, T]. \end{cases} \quad (52)$$

We also give a simple estimation of the diameter of the cell $B_{nkij}(t)$ as follows:

$$\text{diam}(B_{nkij}(t)) < \begin{cases} e^{Lt \frac{\sqrt{2}}{n}} & \text{if } t \in [0, t_{nkij}^0], \\ e^{-C_n(t - t_{nkij}^0)} e^{Lt_{nkij}^0 \frac{\sqrt{2}}{n}} & \text{if } t \in [t_{nkij}^0, T - t_{nkij}^1], \\ e^{-C_n(t_{nkij}^1 - t_{nkij}^0)} e^{LT \frac{\sqrt{2}}{n}} & \text{if } t \in [T - t_{nkij}^1, T], \end{cases} \quad (53)$$

where L is the Lipschitz constant of the vector field v . The estimate in the first interval is a classical application of Gronwall lemma, recalling that $\text{diam}(B_{nkij}^0) < \frac{\sqrt{2}}{n}$. For the estimate in the second interval, we use the fact that the term $v(\hat{x}_{nkij}(t)) + \hat{u}_n(t, \hat{x}_{nkij}(t))$ in w_{nkij} is constant with respect to the space variable, while for the term $C_n(\hat{x}_{nkij}(t) - x)$ we again apply the Gronwall lemma. In the third interval, we again apply the Gronwall lemma and estimate $t \leq T$.

We now prove that $B_{nkij}(t)$ is contained in ω for all $t \in [t_{nkij}^0, T - t_{nkij}^1]$. Indeed, it is sufficient to recall from Step 2.3 that one can choose the trajectory $\hat{x}_{nkij}(t)$ to be contained in ω_n for all $t \in [t_{nkij}^0, T - t_{nkij}^1]$. Now observe that for all $t \in [t_{nkij}^0, T - t_{nkij}^1]$ it holds $\hat{x}_{nkij}(t) \in B_{nkij}(t) \cap \omega_n$, that $\text{diam}(B_{nkij}(t)) < e^{LT \frac{\sqrt{2}}{n}}$ and recall the definition (35) of ω_n with $f_n = e^{LT \frac{\sqrt{2}}{n}}$. Then, $B_{nkij}(t)$ is contained in ω .

We now prove that, given a fixed n and choosing a sufficiently large C_n , for each pair of distinct indexes $nkij, nk'ij' \in I_{n,\varepsilon}^0$ the sets $B_{nkij}(t), B_{nk'ij'}(t)$ are disjoint. Assume for simplicity that it holds $t_{nkij}^0 \leq t_{nk'ij'}^0$. First observe that at time $t = 0$ the sets $B_{nkij}^0, B_{nk'ij'}^0$ are disjoint by construction. We then separate the time interval $[0, T]$ in five intervals:

1. Interval $[0, t_{nkij}^0]$: both sets are displaced via the flow defined by the Lipschitz vector field v , that keeps them disjoint.
2. Interval $[t_{nkij}^0, t_{nk'i'j'}^0]$: observe that the set $B_{nk'i'j'}(t)$ evolves via the flow defined by the Lipschitz vector field v , hence its diameter can grow of a factor $e^{L(t_{nk'i'j'}^0 - t_{nkij}^0)}$. For $t \leq t_{nkij}^1$, by choosing a constant

$$C_n > 2e^{LT} > e^{L(t_{nk'i'j'}^0 - t_{nkij}^0)}$$

sufficiently large, one can reduce the diameter of the set $B_{nkij}(t)$ to have it disjoint with respect to $B_{nk'i'j'}$. For $t > t_{nkij}^1$, it is sufficient to observe that both sets are displaced by the Lipschitz vector field v , then apply Case 1.

3. Interval $[t_{nk'i'j'}^0, \min(t_{nkij}^1, t_{nk'i'j'}^1)]$, when non-empty: recall that the trajectories $\hat{x}_{nkij}(t)$, $\hat{x}_{nk'i'j'}(t)$ are disjoint at each time, since the vector field \hat{u}_n defined in Step 2.3 satisfies the Carathéodory condition. Since in this time interval one can act on both sets with the controls w_{nkij} and $w_{nk'i'j'}$, by choosing C_n sufficiently large one can concentrate the sets in two sufficiently small neighborhoods around disjoint trajectories, then having disjoint sets.
4. Interval $[\min(t_{nkij}^1, t_{nk'i'j'}^1), \max(t_{nkij}^1, t_{nk'i'j'}^1)]$ is equivalent to Case 2.
5. Interval $[\max(t_{nkij}^1, t_{nk'i'j'}^1), T]$ is equivalent to Case 1.

Then, given a fixed n , for each pair of distinct indexes $nkij$, $nk'i'j'$ one can choose a sufficiently large C_n ensuring that the sets $B_{nkij}(t)$, $B_{nk'i'j'}(t)$ are disjoint. By observing that the number of pairs are finite, one can choose a sufficiently large C_n for which the property holds for all pairs of indexes.

We are now ready to define the control u_n approximately steering μ^0 to μ^1 in time $T = S(\mu^0, \mu^1, \omega) + s$. For each time $t \in [0, T]$, first define

$$u_n(t, x) := w_{nkij}(t, x) - v(x) \quad \text{if } t \in [t_{nkij}^0, T - t_{nkij}^1], x \in B_{nkij}(t), \quad (54)$$

where w_{nkij} is defined in (51). It is clear that such definition is well-posed, since the $B_{nkij}(t)$ are disjoint at each time. We now complete the definition of u_n as follows:

$$u_n(t, x) := \begin{cases} w_{nkij}(t, x) - v(x) & \text{for all } t \in [t_{nkij}^0, T - t_{nkij}^1], x \in B_{nkij}(t), \\ \text{Lip. spline} & \text{if } t \in [t_{nkij}^0, T - t_{nkij}^1] \text{ for some } nkij \in I_{n,\varepsilon}^0, \\ \text{in the } x \text{ variable} & \\ 0 & \text{for all } x \in \omega^c, \\ 0 & \text{for all other } t \in [0, T]. \end{cases} \quad (55)$$

Observe that $t \in [t_{nkij}^0, T - t_{nkij}^1]$ and $x \in B_{nkij}(t)$ imply that $B_{nkij} \in \omega$, *i.e.* that the control is allowed to have non-zero value. Also observe that $u_n = 0$ on the boundary of ω , that implies that $\mathbb{1}_\omega u_n$ is also Lipschitz for each time. Regularity with respect to time is ensured by the fact that discontinuities are allowed on a finite number of times t_{nkij}^l only. Thus $\mathbb{1}_\omega u_n$ satisfies the Carathéodory condition, hence it is an admissible control.

We finally observe the following key property: when applying the control $\mathbb{1}_\omega u_n$ to the System (2), the dynamics of the sets B_{nkij}^0 satisfies (52). We will see in the next step that such property is the key to ensure approximate controllability of μ^0 to μ^1 at time T .

Step 3.2: We now prove that the control u_n defined in (54)-(55) provides approximate controllability of μ^0 to μ^1 at time T . Recall that the solution $\mu_n(t)$ to System (2) with control $\mathbb{1}_\omega u_n$ starting from μ^0 is

$$\mu_n(t) := \Phi_t^{v + \mathbb{1}_\omega u_n} \# \mu^0.$$

We aim to prove that the distance $W_1(\mu_n(T), \mu^1)$ is less than ε for n large enough.

Recall the decomposition $\mu^0 = \tilde{\mu}_{n,\varepsilon}^0 + \bar{\mu}_{n,\varepsilon}^0$ introduced in (46), and similarly for μ^1 . One can then write $\mu_n(T) = \Phi_T^{v+1\omega u_n} \# \tilde{\mu}_{n,\varepsilon}^0 + \Phi_T^{v+1\omega u_n} \# \bar{\mu}_{n,\varepsilon}^0$. We estimate

$$W_1(\mu_n(T), \mu^1) \leq W_1(\Phi_T^{v+1\omega u_n} \# \tilde{\mu}_{n,\varepsilon}^0, \tilde{\mu}_{n,\varepsilon}^1) + W_1(\Phi_T^{v+1\omega u_n} \# \bar{\mu}_{n,\varepsilon}^0, \bar{\mu}_{n,\varepsilon}^1). \quad (56)$$

For the first term, recall the decomposition $\tilde{\mu}_{n,\varepsilon}^0 = \sum_{nkij \in I_{n,\varepsilon}^0} \tilde{\mu}_{nkij}^0$ with the supports $\text{supp}(\tilde{\mu}_{nkij}^0) \subset B_{nkij}^0$ by construction. Then, it holds

$$\Phi_T^{v+1\omega u_n} \# \tilde{\mu}_{n,\varepsilon}^0 = \sum_{nkij \in I_{n,\varepsilon}^0} \Phi_T^{v+1\omega u_n} \# \tilde{\mu}_{nkij}^0 \quad (57)$$

and $\text{supp}(\Phi_T^{v+1\omega u_n} \# \tilde{\mu}_{nkij}^0) \subset B_{nkij}(T)$. For each $nkij \in I_{n,\varepsilon}^0$, we provide the estimate

$$\begin{aligned} W_1(\Phi_T^{v+1\omega u_n} \# \tilde{\mu}_{nkij}^0, \tilde{\mu}_{nkij}^1) &\leq W_1\left(\Phi_T^{v+1\omega u_n} \# \tilde{\mu}_{nkij}^0, \left(\frac{1}{n^4} - \frac{1}{n^6}\right) \delta_{x_{nkij}^1}\right) \\ &\quad + W_1\left(\left(\frac{1}{n^4} - \frac{1}{n^6}\right) \delta_{x_{nkij}^1}, \tilde{\mu}_{nkij}^1\right) \\ &\leq \left(\frac{1}{n^4} - \frac{1}{n^6}\right) \text{diam}(B_{nkij}(T)) + \left(\frac{1}{n^4} - \frac{1}{n^6}\right) \text{diam}(B_{nkij}^1) \\ &< \left(\frac{1}{n^4} - \frac{1}{n^6}\right) (e^{LT} + 1) \frac{\sqrt{2}}{n}, \end{aligned} \quad (58)$$

with the following observation: the point $\hat{x}_{nkij}(T) = x_{nkij}^1$ belongs both to $B_{nkij}(T)$ by construction of such set in (52), and to B_{nkij}^1 by construction of such cell in Step 1. Moreover, the mass of both $\tilde{\mu}_{nkij}^0$ and $\tilde{\mu}_{nkij}^1$ is $\frac{1}{n^4} - \frac{1}{n^6}$, then one can apply the triangular inequality (58), then estimate each term by observing that rays in a transference plan sending $\Phi_T^{v+1\omega u_n} \# \tilde{\mu}_{nkij}^0$ to $\left(\frac{1}{n^4} - \frac{1}{n^6}\right) \delta_{x_{nkij}^1}$ have length smaller than $\text{diam}(B_{nkij}(T))$, and similarly for the second term. Estimates of the diameters are consequences of (53) and of the construction of B_{nkij}^1 .

For the second term of (56), first recall that it holds

$$r_{n,\varepsilon} := \bar{\mu}_{n,\varepsilon}^1(\mathbb{R}^d) = 1 - (1 - \varepsilon/4R)(n^4 - n^3)(1/n^4 - 1/n^6) \xrightarrow{n \rightarrow \infty} \varepsilon/4R$$

(see (47) and (49)). Take now any point $x^0 \in \text{supp}(\mu^0)$ and study the trajectory given by the flow $\Phi_t^{v+1\omega u_n}(x^0)$. Consider the set of times $t \in [0, T]$ for which $\Phi_t^{v+1\omega u_n}(x^0) \in \omega$, that is nonempty due to the Geometric Condition 1. Take \bar{t} the supremum of such times and observe that it holds

$$\bar{x} := \Phi_{\bar{t}}^{v+1\omega u_n}(x^0) \in \bar{\omega}.$$

Moreover, the control does not act on the time interval $[\bar{t}, T]$, i.e. $\Phi_t^{v+1\omega u_n}(x^0) = \Phi_t^v(x^0)$. As a consequence, using (48), for n large enough, it holds

$$W_1(\Phi_T^{v+1\omega u_n} \# \bar{\mu}_{n,\varepsilon}^0, \bar{\mu}_{n,\varepsilon}^1) \leq 2R\bar{\mu}_{n,\varepsilon}^1(\mathbb{R}^d) = 2Rr_{n,\varepsilon}. \quad (60)$$

Summing up, from (56)-(57)-(59)-(60) it holds

$$W_1(\mu_n(T), \mu^1) \leq (n^4 - n^3) \left(\frac{1}{n^4} - \frac{1}{n^6}\right) (e^{LT} + 1) \frac{\sqrt{2}}{n} + 2Rr_{n,\varepsilon} \rightarrow \varepsilon/2.$$

For n large enough, $W_1(\mu_n(T), \mu^1)$ is then smaller than ε . Thus System (1) is approximatively controllable for each $T > S(\mu^0, \mu^1, \omega)$.

We now prove **Item (ii)** of Theorem 1.4. To prove it, we first need to recall some results about evolved sets and the Lebesgue measure of their boundaries, which proofs directly follow from the definitions and are then omitted.

Lemma 4.2. *Let ω be an open set and v a Lipschitz vector field. For $t > 0$, define the **evoluted set***

$$\omega^t := \cup_{\tau \in (0,t)} \Phi_\tau^v(\omega).$$

Then, the following statements hold:

1. *If $0 < t_1 < t_2$, then*

$$\omega^{t_1} \subset \omega^{t_2}.$$

2. *For each $t > 0$ it holds $\omega \subset \omega^t$ and $\Phi_t^v(\omega) \subset \omega^t$.*

3. *The set ω^t is open.*

4. *Let μ^0 be a probability measure, with compact support, absolutely continuous with respect to the Lebesgue measure and satisfying the first part of the Geometric Condition 1. Let $x_0 \in \text{supp}(\mu^0)$ and $t_0(x_0)$ the corresponding infimum time to enter ω as defined in (4). Let $\mathbb{1}_\omega u$ be a control satisfying the Carathéodory condition. Then, $t > t_0(x_0)$ implies*

$$\Phi_t^{v+\mathbb{1}_\omega u}(x_0) \in \omega^{t-t_0(x_0)}. \quad (61)$$

Remark 8. The key interest of estimate (61) is that the flow depends on the choice of the control $\mathbb{1}_\omega u$, but that the set $\omega^{t-t_0(x_0)}$ does not depend on it.

We are now ready to prove **Item (ii)** of Theorem 1.4. Consider

$$T \in (S^*(\mu^0, \mu^1), S(\mu^0, \mu^1)].$$

By definition of $S(\mu^0, \mu^1)$, it exists $m \in [0, 1]$ such that

$$T < \mathcal{F}_{\mu^0}^{-1}(m) + \mathcal{B}_{\mu^1}^{-1}(1 - m). \quad (62)$$

Define $\bar{t} := \mathcal{F}_{\mu^0}^{-1}(m)$ and the set

$$A := \{x \in \text{supp}(\mu^0) \text{ s.t. } t_0(x_0) > \bar{t}\}.$$

By definition of $\mathcal{F}_{\mu^0}^{-1}$, it holds $\mu^0(A) = 1 - m$.

Fix now any control $\mathbb{1}_\omega u$. By definition of the quantity $S^*(\mu^0, \mu^1)$, for each $x_0 \in \text{supp}(\mu^0)$, it holds $t_0(x_0) < T$. One can then apply Lemma 4.2, statements 1 and 4, in the two following cases:

- if $x_0 \in \text{supp}(\mu^0) \setminus A$, it holds $\Phi_T^{v+\mathbb{1}_\omega u}(x_0) \in \omega^{T-t_0(x_0)} \subset \omega^T$;
- if $x_0 \in A$, it holds $\Phi_T^{v+\mathbb{1}_\omega u}(x_0) \in \omega^{T-t_0(x_0)} \subset \omega^{T-\bar{t}}$. This implies that

$$\Phi_T^{v+\mathbb{1}_\omega u}(A) \subset \omega^{T-\bar{t}}. \quad (63)$$

Consider now the solution $\mu(t)$ to (2) associated to the control $\mathbb{1}_\omega u$. We already recalled that $\mu(t) = \Phi_t^{v+\mathbb{1}_\omega u} \# \mu^0$. By definition of the push-forward and applying (63), we can compute

$$\mu(T)(\omega^{T-\bar{t}}) = \mu^0 \left((\Phi_T^{v+\mathbb{1}_\omega u})^{-1} \left(\omega^{T-\bar{t}} \right) \right) \geq \mu^0(A) = 1 - m. \quad (64)$$

We now aim to prove that $\mu^1(\omega^{T-\bar{t}}) < 1 - m$. Recall that $\omega^{T-\bar{t}}$ is open, by Lemma 4.2, statement 3. Take now $x_1 \in \omega^{T-\bar{t}} \cap \text{supp}(\mu^1)$, and observe that it holds $x_1 \in \Phi_t^v(\omega)$ for some $t \in (0, T - \bar{t})$. This implies that $\Phi_{-t}^v(x_1) \in \omega$, hence $t_1(x_1) \leq t \leq T - \bar{t}$. Since such property holds for any $x_1 \in \omega^{T-\bar{t}} \cap \text{supp}(\mu^1)$, this

implies that $\mathcal{B}_{\mu^1}(T - \bar{t}) \geq \mu^1(\omega^{T-\bar{t}})$, by definition of \mathcal{B}_{μ^1} . If $\mu^1(\omega^{T-\bar{t}}) \geq 1 - m$, then $\mathcal{B}_{\mu^1}^{-1}(1 - m) \leq T - \bar{t}$, hence $T \geq \bar{t} + \mathcal{B}_{\mu^1}^{-1}(1 - m)$. This contradicts (62), thus it holds $\mu^1(\omega^{T-\bar{t}}) < 1 - m$.

Recall now that $\partial(\omega^{T-\bar{t}})$ has zero Lebesgue measure by the additional hypothesis in Item (ii) of Theorem 1.4. Also recall that μ^1 is absolutely continuous with respect to the Lebesgue measure. Thus, $\mu^1(\overline{\omega^{T-\bar{t}}}) < 1 - m$, hence by standard approximation of measurable sets, there exists an open set $\mathcal{O} \supset \overline{\omega^{T-\bar{t}}}$ such that $\mu^1(\mathcal{O}) < 1 - m$. Observe that such set depends on μ^1 only, thus it does not depend on the choice of the control u . Also observe that, by a compactness argument, we have that

$$D := \inf\{\|x - y\| \text{ s.t. } x \in \overline{\omega^{T-\bar{t}}}, y \notin \mathcal{O}\}$$

is strictly positive.

We now prove that for each control $\mathbb{1}_\omega u$, the associated solution $\mu(t)$ to (2) satisfies

$$W_1(\mu(T), \mu^1) \geq D(1 - m - \mu^1(\mathcal{O})). \quad (65)$$

The idea is to observe that the $1 - m$ mass of $\mu(T)$ in $\omega^{T-\bar{t}}$ cannot be completely transferred to the set \mathcal{O} , since $\mu^1(\mathcal{O}) < 1 - m$. Thus, a part of it needs to be transferred to $\mathbb{R}^d \setminus \mathcal{O}$, for which the distance of transfer is larger than D . More formally, consider a transference plan $\pi \in \Pi(\mu(T), \mu^1)$: recall that $\pi(\mathbb{R}^d \times \mathcal{O}) = \mu^1(\mathcal{O}) < 1 - m$, thus $\pi(\omega^{T-\bar{t}} \times \mathcal{O}) \leq \mu^1(\mathcal{O}) < 1 - m$, while $\pi(\omega^{T-\bar{t}} \times \mathbb{R}^d) = \mu(T)(\omega^{T-\bar{t}}) \geq 1 - m$ by (64). Thus it holds

$$\pi(\omega^{T-\bar{t}} \times (\mathbb{R}^d \setminus \mathcal{O})) \geq 1 - m - \mu^1(\mathcal{O}).$$

Observe that $(x, y) \in \omega^{T-\bar{t}} \times (\mathbb{R}^d \setminus \mathcal{O})$ implies $\|x - y\| \geq D$. As a consequence, it holds

$$\int_{\mathbb{R}^d \times \mathbb{R}^d} \|x - y\| d\pi(x, y) \geq \int_{\omega^{T-\bar{t}} \times (\mathbb{R}^d \setminus \mathcal{O})} D d\pi(x, y) = D(1 - m - \mu^1(\mathcal{O})).$$

Passing to the infimum among all transference plans $\pi \in \Pi(\mu(T), \mu^1)$, it holds (65). Observe that neither D nor $1 - m - \mu^1(\mathcal{O})$ depend on the choice of the control $\mathbb{1}_\omega u$, thus there exists no sequence $\mathbb{1}_\omega u_n$ such that the associated solution μ_n satisfies $W_1(\mu_n(T), \mu^1) \rightarrow 0$. Hence, the system is not approximately controllable at time T . \square

Remark 9. We give now two examples in which System (1) is never exactly controllable on $[0, S^*(\mu^0, \mu^1))$ and another where System (1) is exactly controllable at each time $T \in [0, S^*(\mu^0, \mu^1))$.

Figure 7 (left). Consider $\omega := (-1, 1) \times (-1.5, 1.5)$. The vector field v is $(1, 0)$, thus uncontrolled trajectories are right translations. Define

$$\begin{cases} \mu^0 := \mathbb{1}_{(-2.5, -2) \times (-1, 1)} dx, \\ \mu^1 := \mathbb{1}_{(2, 2.5) \times (-1, 1)} dx. \end{cases}$$

The time $S^*(\mu^0, \mu^1)$ at which we can act on the particles and the minimal time $S(\mu^0, \mu^1)$ are respectively equal to 1.5 and 2.5. We observe that for each time $T \in [0, S^*(\mu^0, \mu^1)]$ System (2) is not approximately controllable. Indeed, each point takes a time $t > 2$ to go from $\text{supp}(\mu^0)$ to $\text{supp}(\mu^1)$, hence one cannot expect approximate controllability for smaller times.

Figure 7 (right). Consider $\omega := (-1, 1) \times (-1.5, 1.5)$. The vector field v is $(-y, x)$, thus uncontrolled trajectories are rotations with constant angular velocity. Define

$$\mu^0 = \mu^1 := \mathbb{1}_{B_1(1,0) \setminus B_{0.5}(1,0)} dx.$$

In this case, both quantities $S^*(\mu^0, \mu^1)$ and $S(\mu^0, \mu^1)$ are equal to π . Since $\Phi_t^v \# \mu^0 = \mu^1$ for all $t \geq 0$, we remark that System (1) is exactly controllable for all $T \in [0, S^*(\mu^0, \mu^1))$.




Figure 7: Left : The macroscopic system is not approximately controllable for each $T \in [0, S^*(\mu^0, \mu^1))$. Right : The macroscopic system is approximately controllable for each $T \in [0, S^*(\mu^0, \mu^1))$.

5 Numerical simulations

In this section, we give some numerical examples of the algorithm developed in the proof of Theorem 1.4 to compute the infimum time and the solution associated to the minimal time problem to approximately steer an AC measure to another. We use a Lagrangian scheme, introduced in [42, 44], for simulations of transport equations.

We construct the mesh in Algorithm 1. We then compute the minimal time in Algorithm 2.

We first give Algorithm 1. To simplify the notations, we assume that the space dimension is $d = 2$.

Algorithm 1 Construction of the meshes \mathcal{T}_n^0 and \mathcal{T}_n^1

Let $n \in \mathbb{N}^*$ and two AC measures μ^0, μ^1 be given. Choose α_0, α_1 such that $\text{supp}(\mu^0), \text{supp}(\mu^1) \subset (\alpha_0, \alpha_1)^2$.

Step 1: Construction, for all $k := (k_1, k_2) \in \{0, \dots, n-1\}^2$, of the sets

$$C_{n,k} := \left[\alpha_0 + (\alpha_1 - \alpha_0) \frac{k_1}{n}, \alpha_0 + (\alpha_1 - \alpha_0) \frac{k_1+1}{n} \right] \times \left[\alpha_0 + (\alpha_1 - \alpha_0) \frac{k_2}{n}, \alpha_0 + (\alpha_1 - \alpha_0) \frac{k_2+1}{n} \right].$$

Step 2: Partition of $C_{n,k}$ into subsets $\{C_{nki}^0\}_i$ with $C_{nki}^0 = [a_i^0, a_{i+1}^0] \times (\alpha_0, \alpha_1)$ such that $\mu_{C_{n,k}}^0(C_{nki}^0) = 1/n^3$ (if $\mu_{C_{n,k}}^0(C_{nki}^0) < 1/n^3$, then we do not partition $C_{n,k}$) and, for each i , partition of C_{nki}^0 into some subsets $\{C_{nki j}^0\}_j$ with $C_{nki j}^0 = [a_i^0, a_{i+1}^0] \times [a_{i(j+1)}^0, a_{i(j+1)+1}^0]$ such that

$$\mu^0(C_{nki j}^0) = 1/n^4.$$

See Figure 5. We define similarly the cells $C_{nki j}^1$.

Step 3: Construction of $B_{nki j}^0 := [b_i^{0-}, b_i^{0+}] \times [b_{i j}^{0-}, b_{i j}^{0+}] \subset C_{nki j}^0$ and $B_{nki j}^1 := [b_i^{1-}, b_i^{1+}] \times [b_{i j}^{1-}, b_{i j}^{1+}] \subset C_{nki j}^1$ such that

$$\mu^0(B_{nki j}^0) = \mu^1(B_{nki j}^1) = \frac{1}{n^4} - \frac{1}{n^6}.$$

See Figure 6.

Step 4: Definition of $\mathcal{T}_n^0 := \bigcup_{kij \in I_n^0} B_{nki j}^0$ and $\mathcal{T}_n^1 := \bigcup_{kij \in I_n^1} B_{nki j}^1$, where, for $l = 0, 1$, I_n^l is the set of kij such that $B_{nki j}^l$ is well defined.

We now recall the algorithm to compute the minimal time. We will assume that ω is convex in order to use the constructive proof of Proposition 7, *i.e.* the agents cross the control area following straight trajectories.

Algorithm 2 Approximate controllability

Let μ^0 and μ^1 be two AC measures satisfying the Geometric Condition 1.

Step 1: Construction of the meshes $\mathcal{T}_n^0 := \bigcup_{kij \in I_n^0} B_{nkij}^0$ and $\mathcal{T}_n^1 := \bigcup_{kij \in I_n^1} B_{nkij}^1$ following Algorithm 1.

Step 2: Definition of $X^l := \{x_{kij}^l : kij \in I_n^l\}$ ($l = 0, 1$) with x_{nkij}^l being a point of $B_{nkij}^l \cap \text{supp}(\mu^l)$

Step 3: Definition of

$$\omega_n := \{x \in \mathbb{R}^d : d(x, \omega^c) > e^{LT} \sqrt{2}/n\}.$$

For $t_{nkij}^l := t^l(x_{nkij}^l, \omega_n)$ with $l = 0, 1$, computation of the minimal time to steer X^0 to X^1 up to a mass $\varepsilon/4R$

$$M_{e, \varepsilon/4R}(X^0, X^1, \gamma) := \max_{1 \leq i \leq n-R} \{t_{nkij}^0 + t_{nkij+R}^1\},$$

where $R := \lfloor n\varepsilon\gamma/4R \rfloor$ and the sequences $\{t_{nkij}^0\}_{kij}$, $\{t_{nkij}^1\}_{kij}$ are increasingly and decreasingly ordered, respectively

Step 4: Computation of the optimal permutations σ_0 and σ_1 minimizing (34) to steer X^0 to X^1 up to a mass $\varepsilon/4R$.

Step 5: Concentration of the mass of B_{nkij}^0 around x_{nkij}^0 in order to obtain no intersection of the cells when they follow the trajectories of $\delta_{x_{nkij}^0}$

Step 6: Computation of the control u_n and the solution μ_n to (2) on $(0, T)$.

5.1 Example 1: the 1-D case

Consider the initial data μ^0 and the target μ^1 defined by

$$\begin{cases} \mu^0 := 0.5 \times \mathbb{1}_{(0,2)} dx, \\ \mu^1 := 0.5 \times \mathbb{1}_{(7,8) \cup (10,11)} dx. \end{cases}$$

Let the velocity field $v := 1$ and the control region $\omega := (5, 6)$ be given. This situation is illustrated in Figure 8. In this case, the infimum time $T_a(\mu^0, \mu^1)$ is 8, which is computed in Step 3 of Algorithm 2. One cannot




Figure 8: Control set ω , initial configuration μ^0 and final one μ^1 for Example 1.

achieve approximate control at such time, but we aim to control the system at time $T = T_a(\mu^0, \mu^1) + \delta$, with $\delta := 0.1$, hence $T = 8.1$. Following Algorithm 2, we obtain the solution presented in Figure 9. The maximal density for this solution is equal to 1.1. It is due to the fact that we concentrate a part of the mass coming from the set $\{x < 5\}$ in the control set, to slow it down. This increase of the maximal density can be seen as a drawback of the method for several key applications, namely for egress problems. Indeed, high concentrations need to be carefully avoided in such settings, since they might induce death by suffocation, that is among the main causes of fatalities in stampedes/crashes (see for instance [30]).

For this reason, in the future we plan to study new control strategies for minimal time problems, in which a constraint on the maximal density is added. Alternatively, we aim to estimate the maximal density value that is reached with optimal strategies.




Figure 9: Example 1: solution at time $t = 0$, $t = 3.4$, $t = 4.6$ and $t = T = 8.1$.




Figure 10: Control set ω , initial configuration μ^0 and final one μ^1 for Example 2.

5.2 Example 2: the 2D case

We now give an example in the 2D case. Consider the initial data μ^0 and the target μ^1 defined by

$$\begin{cases} \mu^0 := \frac{1}{8} \times \mathbb{1}_{(0,4) \times (1,3)} dx, \\ \mu^1 := \frac{1}{16} \times \mathbb{1}_{(8,14) \times (0,4) \setminus (9,13) \times (1,3)} dx. \end{cases}$$

We fix the velocity field $v := (1, 0)$ and the control region $\omega := (5, 7) \times (0, 4)$. This situation is illustrated in Figure 10. Again, in this case $T_a(\mu^0, \mu^1) = 8$. Since it is not possible to approximately steer μ^0 to μ^1 at such time, we control the system at time $T = T_a(\mu^0, \mu^1) + \delta$, with $\delta := 1$, hence $T = 9$. Following Algorithm 2, we present the solution in Figure 11. As in the previous example, we observe a high concentration of the crowd in the control region, in this case with a maximal density equal to 8.

A Proof of Lemma 1.5

In this section, we prove Lemma 1.5. We first give the definition of the uniform interior cone condition and state a related result.




Figure 11: Example 2: solution at time $t = 0$, $t = 3$, $t = 6$ and $t = T = 9$.

Definition A.1. Let

$$C(x_0, n, \alpha, r) := \left\{ x \in B_r(x_0) \text{ s.t. } \begin{aligned} n \cdot (x - x_0) &> 0, \\ \|x - x_0 - n(n \cdot (x - x_0))\| &< \alpha n \cdot (x - x_0) \end{aligned} \right\}$$

be a (open) cone centered in x_0 , with normal unit vector n , tangent α and radius r .

We say that a set A satisfies the uniform interior cone condition if there exist uniform α, r such that for all $x_0 \in \partial A$ there exists n such that

$$C(x_0, n, \alpha, r) \subset A.$$

Lemma A.2. Let v be a C^1 vector field, and A an open bounded set satisfying the uniform interior cone condition. Then, the following statements hold.

1. The set $\Phi_t^v(A)$ satisfies the uniform interior cone condition;
2. For each $t > 0$, the set $A^t = \cup_{\tau \in (0, t)} \Phi_\tau^v(A)$ satisfies the uniform interior cone condition.

The proof is omitted. It can be easily recovered by a first-order expansion of the flow, in which parameters depend continuously on the point, since v is chosen to be C^1 .

We are now ready to prove Lemma 1.5. Since ω satisfies the uniform interior cone condition, by Lemma A.2, Statement 2, it holds that ω^t satisfies the uniform interior cone condition, with uniform parameters α', r' . We now prove that this implies that $\partial(\omega^t)$ has zero Lebesgue measure. Since ω^t is open and bounded, then it is measurable, thus its characteristic function $\mathbb{1}_{\omega^t}$ is in $L^1(\mathbb{R}^d)$. It is now sufficient to prove that no point of the boundary is a Lebesgue point with respect to the Lebesgue measure. Indeed, observe from one side that for each $x \in \partial(\omega^t)$ it holds $\mathbb{1}_{\omega^t}(x) = 0$ since ω^t is open. On the other side, for $r < r'$ it holds

$$\frac{1}{\mathcal{L}(B_r(x))} \int_{B_r(x)} \mathbb{1}_{\omega^t}(y) d\mathcal{L}(y) \geq \frac{\mathcal{L}(C(x, m, \alpha', r') \cap B_r(x))}{\mathcal{L}(B_r(x))}.$$

By a simple dilation and rototranslation argument, we remark that such term coincides with $\frac{\mathcal{L}(C(0, m, \alpha', 1))}{\mathcal{L}(B_1(0))}$, that is a strictly positive constant, independent on r . As a consequence, it holds

$$0 = \mathbb{1}_{\omega^t}(x) \neq \lim_{r \rightarrow 0} \frac{1}{\mathcal{L}(B_r(x))} \int_{B_r(x)} \mathbb{1}_{\omega^t}(y) d\mathcal{L}(y),$$

hence x is not a Lebesgue point. Since $x \in \partial(\omega^t)$ is generic, then no point in $\partial(\omega^t)$ is a Lebesgue point. Since the set of points that are not Lebesgue points for a measurable function has zero Lebesgue measure (Lebesgue-Besicovitch differentiation theorem, see *e.g.* [25, Sec. 1.7]), then $\partial(\omega^t)$ has zero Lebesgue measure.

References

- [1] Y. Achdou and M. Laurière, Mean field type control with congestion, *Applied Mathematics & Optimization*, **73** (2016), 393–418.
- [2] Y. Achdou, F. Camilli and I. Capuzzo-Dolcetta, Mean field games: Numerical methods for the planning problem, *SIAM Journal on Control and Optimization. on Contr. and Opt.*, **50** (2012), 77–109.
- [3] Y. Achdou and M. Laurière, On the system of partial differential equations arising in mean field type control, *Discrete Contin. Dyn. Syst.*, **35** (2015), 3879–3900.
- [4] A. A. Agrachev and Y. Sachkov, *Control theory from the geometric viewpoint*, vol. 87, Springer Science & Business Media, 2013.
- [5] R. Axelrod, *The Evolution of Cooperation: Revised Edition*, Basic Books, 2006.
- [6] N. Bellomo, P. Degond and E. Tadmor, *Active Particles, Volume 1: Advances in Theory, Models, and Applications*, Modeling and Simulation in Science, Engineering and Technology, Springer, 2017.
- [7] B. Bonnet, A Pontryagin Maximum Principle in Wasserstein Spaces for Constrained Optimal Control Problems, *ESAIM: COCV*, to appear.
- [8] B. Bonnet and F. Rossi, The Pontryagin Maximum Principle in the Wasserstein Space, *Calc. Var. PDE*, **58:11** (2019).
- [9] A. Bressan and B. Piccoli, *Introduction to the mathematical theory of control*, AIMS, 2007.
- [10] F. Bullo, J. Cortés and S. Martínez, *Distributed Control of Robotic Networks: A Mathematical Approach to Motion Coordination Algorithms*, Princeton University Press, 2009.
- [11] C. Calcaterra and A. Boldt, Lipschitz Flow-box Theorem, *J. Math. An. Appl.*, **338** (2008), 1108 - 1115.
- [12] S. Camazine, *Self-organization in Biological Systems*, Princeton studies in complexity, Princeton University Press, 2003.
- [13] C. Canudas-de-Wit, L. L. Ojeda and A. Y. Kibangou, Graph constrained-CTM observer design for the Grenoble south ring, *IFAC Proceedings Volumes*, **45** (2012), 197–202.
- [14] M. Caponigro, B. Piccoli, F. Rossi and E. Trélat, Mean-field sparse Jurdjevic-Quinn control, *Math. Models Methods Appl. Sci.*, **27** (2017), 1223–1253.
- [15] M. Caponigro, B. Piccoli, F. Rossi and E. Trélat, Sparse Jurdjevic-Quinn stabilization of dissipative systems, *Automatica J. IFAC*, **86** (2017), 110–120.
- [16] R. Carmona, F. Delarue and A. Lachapelle, Control of McKean–Vlasov dynamics versus mean field games, *Mathematics and Financial Economics*, **7** (2013), 131–166.
- [17] G. Cavagnari, Regularity results for a time-optimal control problem in the space of probability measures, *Mathematical Control and Related Fields*, **7** (2017), 213–233.
- [18] G. Cavagnari, A. Marigonda and B. Piccoli, Optimal synchronization problem for a multi-agent system, *Networks and Heterogeneous Media*, **12** (2017), 277–295.
- [19] G. Cavagnari, A. Marigonda and B. Piccoli, Averaged time-optimal control problem in the space of positive Borel measures, *ESAIM: Contr., Opt. and Cal. of Var.*, **24** (2018), 721–740.

- [20] T. Champion, L. De Pascale and P. Juutinen, The ∞ -Wasserstein distance: local solutions and existence of optimal transport maps, *SIAM J. Math. Anal.*, **40** (2008), 1–20.
- [21] J.-M. Coron, *Control and nonlinearity*, vol. 136 of Mathematical Surveys and Monographs, American Mathematical Society, Providence, RI, 2007.
- [22] E. Cristiani, B. Piccoli and A. Tosin, Multiscale modeling of pedestrian dynamics, 2014.
- [23] F. Cucker and S. Smale, Emergent behavior in flocks, *IEEE Trans. Automat. Control*, **52** (2007), 852–862.
- [24] M. Duprez, M. Morancey and F. Rossi, Approximate and exact controllability of the continuity equation with a localized vector field, *SIAM J. Control Optim.*, **57** (2019), 1284–1311.
- [25] L. Evans and R. Gariepy, *Measure theory and fine properties of functions*, Revised edition, Textbooks in Mathematics, CRC Press, Boca Raton, FL, 2015.
- [26] A. Ferscha and K. Zia, Lifebelt: Crowd evacuation based on vibro-tactile guidance, *IEEE Pervasive Computing*, **9** (2010), 33–42.
- [27] M. Fornasier and F. Solombrino, Mean-field optimal control, *ESAIM: Control, Optimisation and Calculus of Variations*, **20** (2014), 1123–1152.
- [28] A. Hegyi, S. Hoogendoorn, M. Schreuder, H. Stoelhorst and F. Viti, Specialist: A dynamic speed limit control algorithm based on shock wave theory, in *Intel. Transp. Syst., 2008. ITSC 2008. 11th Intern. IEEE Conf. on*, IEEE, 2008, 827–832.
- [29] D. Helbing and R. Calek, *Quantitative Sociodynamics: Stochastic Methods and Models of Social Interaction Processes*, Theo. and Dec. Lib. B, Springer Neth., 2013.
- [30] D. Helbing, I. J. Farkas and T. Vicsek, Crowd disasters and simulation of panic situations, in *The Science of Disasters*, Springer, 2002, 330–350.
- [31] M. Jackson, *Social and Economic Networks*, Princeton University Press, 2010.
- [32] P. Jameson Graber, A. R. Mészáros, F. J. Silva, D. Tonon, The planning problem in Mean Field Games as regularized mass transport, arXiv:1811.02706.
- [33] V. Jurdjevic, *Geometric control theory*, vol. 52, Cambridge university press, 1997.
- [34] M. Krein and D. Milman, On extreme points of regular convex sets, *Studia Math.*, **9** (1940), 133–138.
- [35] V. Kumar, N. Leonard and A. Morse, *Cooperative Control: A Post-Workshop Volume, 2003 Block Island Workshop on Cooperative Control*, Lecture Notes in Control and Information Sciences, Springer Berlin Heidelberg, 2004.
- [36] Z. Lin, W. Ding, G. Yan, C. Yu and A. Giua, Leader–follower formation via complex Laplacian, *Automatica*, **49** (2013), 1900 – 1906.
- [37] J. Lohéac, E. Trélat and E. Zuazua, Minimal controllability time for the heat equation under unilateral state or control constraints, *Math. Models Methods Appl. Sci.*, **27** (2017), 1587 – 1644.
- [38] P. B. Luh, C. T. Wilkie, S.-C. Chang, K. L. Marsh and N. Olderman, Modeling and optimization of building emergency evacuation considering blocking effects on crowd movement, *IEEE Transactions on Automation Science and Engineering*, **9** (2012), 687–700.
- [39] S. Motsch and E. Tadmor, Heterophilious dynamics enhances consensus, *SIAM Review*, **56** (2014), 577–621.

- [40] C. Orrieri, A. Porretta, G. Savaré, A variational approach to the mean field planning problem, *Journal of Functional Analysis*, to appear.
- [41] B. Piccoli and F. Rossi, On properties of the Generalized Wasserstein distance, *Arch. Ration. Mech. Anal.*, **222** (2016), 1339–1365.
- [42] B. Piccoli and F. Rossi, Transport equation with nonlocal velocity in Wasserstein spaces: convergence of numerical schemes, *Acta Appl. Math.*, **124** (2013), 73–105.
- [43] B. Piccoli, F. Rossi and E. Trélat, Control to flocking of the kinetic Cucker-Smale model, *SIAM J. Math. Anal.*, **47** (2015), 4685–4719.
- [44] B. Piccoli and A. Tosin, Time-evolving measures and macroscopic modeling of pedestrian flow, *Archive for Rational Mechanics and Analysis*, **199** (2011), 707–738.
- [45] A. Porretta, On the planning problem for a class of mean field games, *Comptes Rendus Mathématique*, **351** (2013), 457–462.
- [46] A. Porretta, On the planning problem for the mean field games system, *Dynamic Games and Applications*, **4** (2014), 231–256.
- [47] R. Sepulchre, Consensus on nonlinear spaces, *Ann. rev. in contr.*, **35** (2011), 56–64.
- [48] E. D. Sontag, *Mathematical control theory: deterministic finite dimensional systems*, vol. 6, Springer Science & Business Media, 2013.
- [49] C. Villani, *Topics in optimal transportation*, vol. 58 of Graduate Studies in Mathematics, American Mathematical Society, Providence, RI, 2003.